


# Learning Outcome: Interference of light and Coherence property of light

8/6/2018



$M \rightarrow \# \text{ of slits}$

$\phi \rightarrow \text{phase difference}$   
from successive slits

Wave component

$$U_m = \sqrt{I_0} \exp[i(m-1)\phi]$$

where  $m=1, 2, \dots, M$

??

Observation  
plane




Total Wave amplitude  $U = \sqrt{I_0} (1 + h + h^2 + \dots + h^{M-1})$  where  $h = e^{j\phi}$

$$= \sqrt{I_0} \cdot \frac{1 - h^M}{1 - h} = \sqrt{I_0} \cdot \frac{1 - e^{jM\phi}}{1 - e^{j\phi}}$$

Total Intensity

$$I = |U|^2 = I_0 \cdot \left| \frac{1 - e^{jM\phi}}{1 - e^{j\phi}} \right|^2 = I_0 \left| \frac{e^{-jM\phi/2} - e^{jM\phi/2}}{e^{-j\phi/2} - e^{j\phi/2}} \right|^2$$

$$I = I_0 \cdot \frac{\sin^2(M\phi/2)}{\sin^2(\phi/2)}$$



$$\frac{M\phi_{min}}{2} = \pi$$

$$\phi_{min} = \frac{2\pi}{M}$$


More # of interfering sources

$\Rightarrow$  Narrower spectral selectivity




# Interferometers

## Common Path



Sagnac

Fabry-Pérot



$$\Delta\phi = \frac{2\pi}{\lambda} n \cdot 2d$$


$$= 2\pi m$$

$n=1$ ,


$$d = m \cdot \frac{\lambda}{2}$$

## Differential Path

Mach-Zehnder



Michelson



$$\phi_2 - \phi_1 = \frac{2\pi}{\lambda} 2n_2 d_2 - \frac{2\pi}{\lambda} 2n_1 d_1$$

$$= \frac{2\pi}{\lambda} 2(n_2 d_2 - n_1 d_1)$$

Optical path length difference

Constructive Interference,  $\phi_2 - \phi_1 = 2\pi m$

If  $n_1 = n_2 = 1$ ,  $d_1 - d_2 = \frac{m\lambda}{2}$


Total Wave amplitude  $U = \sqrt{I_0} (1 + h + h^2 + \dots + h^{M-1})$  where  $h = e^{j\phi}$

$$= \sqrt{I_0} \cdot \frac{1 - h^M}{1 - h} = \sqrt{I_0} \cdot \frac{1 - e^{jM\phi}}{1 - e^{j\phi}}$$

Total Intensity

$$I = |U|^2 = I_0 \cdot \left| \frac{1 - e^{jM\phi}}{1 - e^{j\phi}} \right|^2 = I_0 \cdot \left| \frac{e^{-jM\phi/2} - e^{jM\phi/2}}{e^{-j\phi/2} - e^{j\phi/2}} \right|^2$$

$$I = I_0 \cdot \frac{\sin^2(M\phi/2)}{\sin^2(\phi/2)}$$



$$M\phi_{min} = \pi$$

$$\phi_{min} = \frac{2\pi}{M}$$

More # of interfering sources

$\Rightarrow$  Narrower spectral selectivity

# of slits  
 $\phi$  = phase difference  
for adjacent slits

# of slits  
 $\phi$  = phase difference  
for adjacent slits

??

