


Lo: Identify the fundamental principles of laser & quantify their characteristics

9/17/2018



Laser oscillation condition

$$E_0 e^{+\gamma/2 \cdot 2L} e^{-\alpha_{int}/2 \cdot 2L} \sqrt{R_1 R_2} \cdot e^{-j k_0 n \cdot 2L} = E_0$$

$$e^{\gamma L} e^{-\alpha_{int} L} \sqrt{R_1 R_2} = 1$$

$$k_0 \cdot n \cdot 2L = 2\pi m$$

①

$\Rightarrow$


$$\gamma = \alpha_{int} + \frac{1}{2L} \ln\left(\frac{1}{R_1 R_2}\right) = \alpha_r$$

②

$\Rightarrow$

$$\frac{2\pi}{\lambda} \cdot n \cdot 2L = 2\pi m$$



$$\gamma = m \cdot \frac{c}{2nL}$$




Avg. Photon lifetime,  $\tau_{ph} \propto \frac{1}{\alpha_r}$

$$\tau_{ph} = \frac{1}{c \cdot \alpha_r} \Rightarrow \alpha_r = \frac{1}{c \cdot \tau_{ph}}$$

$$\Rightarrow N_{th} = \frac{1}{c \cdot \tau_{ph} \cdot \sigma}$$






①  $\Rightarrow$

$$\gamma = \alpha_{int} + \frac{1}{2L} \ln\left(\frac{1}{R_1 R_2}\right) = \alpha_r$$

②  $\Rightarrow$

$$\frac{2\pi}{\lambda} \cdot n \cdot 2L = 2\pi m$$


$$\nu = m \cdot \frac{c}{2nL}$$



For laser oscillation,  $\gamma > \alpha_r$

Assume  $\sigma_e = \sigma_a = \sigma$ ,  $\sigma N > \alpha_r$

Threshold inversion,  $N_m = \frac{\alpha_r}{\sigma}$



$$\frac{\gamma_o(\nu)}{1 + \phi_s(\nu)/\phi_{sat}} = \alpha_r \text{ at threshold}$$