

Module 6: Dimensionality Reduction

Lecture 28: Dimensionality Reduction

Prev topic

Next topic

Prev page

Next page

The Lecture Contains:

- General idea
- Distortion
- Reduction to one dimension

Module 6: Dimensionality Reduction

Lecture 28: Dimensionality Reduction

[Prev topic](#)[Next topic](#)[Prev page](#)[Next page](#)

General idea

- Feature vector \boldsymbol{v} in k dimensions
- Transformation f on to yield another feature vector $f(\boldsymbol{v})$ of dimensionality $k' < k$
- Distance measure may change from d to d'

Module 6: Dimensionality Reduction

Lecture 28: Dimensionality Reduction

[Prev topic](#)[Next topic](#)[Prev page](#)[Next page](#)

General idea

- Feature vector v in k dimensions
- Transformation f on to yield another feature vector $f(v)$ of dimensionality $k' < k$
- Distance measure may change from d to d'
- Aim is to maintain, as far as possible

$$d(x, y) \approx d'(f(x), f(y))$$

Module 6: Dimensionality Reduction

Lecture 28: Dimensionality Reduction

Prev topic

Next topic

Prev page

Next page

General idea

- Feature vector v in k dimensions
- Transformation f on to yield another feature vector $f(v)$ of dimensionality $k' < k$
- Distance measure may change from d to d'
- Aim is to maintain, as far as possible

$$d(x, y) \approx d'(f(x), f(y))$$

- Pruning property

$$\forall x, y, d'(f(x), f(y)) \leq d(x, y)$$

- Mappings that obey pruning property are **contractive**
- Useful for range queries

Module 6: Dimensionality Reduction

Lecture 28: Dimensionality Reduction

Prev topic

Next topic

Prev page

Next page

General idea

- Feature vector v in k dimensions
- Transformation f on to yield another feature vector $f(v)$ of dimensionality $k' < k$
- Distance measure may change from d to d'
- Aim is to maintain, as far as possible

$$d(x, y) \approx d'(f(x), f(y))$$

- Pruning property

$$\forall x, y, d'(f(x), f(y)) \leq d(x, y)$$

- Mappings that obey pruning property are **contractive**
- Useful for range queries
- Proximity-preserving property

$$\forall x, y, z, d(x, y) \leq d(x, z) \implies d'(f(x), f(y)) \leq d'(f(x), f(z))$$

- Useful for nearest-neighbor queries

Module 6: Dimensionality Reduction

Lecture 28: Dimensionality Reduction

[Prev topic](#)[Next topic](#)[Prev page](#)[Next page](#)

Distortion

$$\frac{1}{c_1} \cdot d(x, y) \leq d'(f(x), f(y)) \leq c_2 \cdot d(x, y)$$

- $c_1, c_2 \geq 1$ provides bounds on distance after dimensionality reduction
- c_1 is contraction and c_2 is expansion

Module 6: Dimensionality Reduction

Lecture 28: Dimensionality Reduction

[Prev topic](#)
[Next topic](#)
[Prev page](#)
[Next page](#)

Distortion

$$\frac{1}{c_1} \cdot d(x, y) \leq d'(f(x), f(y)) \leq c_2 \cdot d(x, y)$$

- $c_1, c_2 \geq 1$ provides bounds on distance after dimensionality reduction
- c_1 is contraction and c_2 is expansion

$$\text{Distortion} = c_1 \cdot c_2$$

- For arbitrary dimensionality reductions,

$$\text{Distortion} = \max_{\forall x, y} \left\{ \frac{d(x, y)}{d'(f(x), f(y))} \right\} \times \max_{\forall x, y} \left\{ \frac{d'(f(x), f(y))}{d(x, y)} \right\}$$

- Contractive embeddings: $c_2 = 1$
- Isometric embeddings: $c_1 = c_2 = 1$

Module 6: Dimensionality Reduction

Lecture 28: Dimensionality Reduction

[Prev topic](#)[Next topic](#)[Prev page](#)[Next page](#)

Reduction to one dimension

- Hashing
- Space-filling curves
- Choosing single dimension
 - Choose dimension with largest spread or largest variance
- Pruning and proximity-preserving properties are generally violated
- Loss of information and distortion are too high
- In general, not very useful