

* CONDITIONAL

ENTROPY

X, Y

Entropy of Y conditioned on X

$$H(Y|X) = \sum_{i=0}^{M-1} P(X=s_i) \cdot H(Y|X=s_i)$$

$$= \sum_{i=0}^{M-1} \Pr(X=s_i) \sum_{j=0}^{N-1} \Pr(Y=\tau_j | X=s_i) \times \log_2 \frac{1}{\Pr(Y=\tau_j | X=s_i)}$$

$$H(Y|X=s_i)$$

$$= \sum_{i=0}^{M-1} \Pr(s_i) \sum_{j=0}^{N-1} \Pr(\tau_j | s_i) \cdot \log_2 \frac{1}{\Pr(\tau_j | s_i)}$$

$$= \sum_{i=0}^{M-1} \sum_{j=0}^{N-1} \frac{\Pr(s_i, \tau_j)}{\Pr(s_i) \Pr(\tau_j | s_i)} \cdot \log_2 \left(\frac{1}{\Pr(\tau_j | s_i)} \right)$$

$$= \sum_{i=0}^{M-1} \sum_{j=0}^{N-1} \Pr(x_i, y_j) \log_2 \frac{1}{\Pr(y_j | x_i)}$$

$H(Y|X)$
 Conditional Entropy
 of Y given X

$H(Y|X)$ = uncertainty
 remaining in Y
 having observed X

$$H(Y|X) + H(X)$$

$$= \sum_{i=0}^{M-1} \sum_{j=0}^{N-1} P_r(\delta_i, \tau_j) \log_2 \frac{1}{P_r(\tau_j | \delta_i)}$$

$H(Y|X)$

$$+ \sum_{i=0}^{M-1} P_r(\delta_i) \log_2 \frac{1}{P_r(\delta_i)}$$

$$\sum_{j=0}^{N-1} \sum_{i=0}^{M-1} P_r(\tau_j, \delta_i) \log_2 \left(\frac{1}{P_r(\delta_i)} \right)$$

$H(X)$

$$\Rightarrow H(Y|X) + H(X)$$

$$= \sum_{i=0}^{M-1} \sum_{j=0}^{N-1} \text{Pr}(\delta_i, \tau_j) \log_2 \frac{1}{\text{Pr}(\tau_j | \delta_i)}$$

$$+ \sum_{i=0}^{M-1} \sum_{j=0}^{N-1} \text{Pr}(\delta_i, \tau_j) \log_2 \frac{1}{\text{Pr}(\delta_i)}$$

$$= \sum_{i=0}^{M-1} \sum_{j=0}^{N-1} \text{Pr}(\delta_i, \tau_j) \log_2 \left(\frac{1}{\underbrace{\text{Pr}(\tau_j | \delta_i) \text{Pr}(\delta_i)}_{= \text{Pr}(\delta_i, \tau_j)}} \right)$$

$$= \sum_{i=0}^{M-1} \sum_{j=0}^{N-1} P_r(s_i, t_j) \log_2 \left(\frac{1}{P_r(s_i, t_j)} \right)$$

$H(X, Y)$

uncertainty in Y given X uncertainty in X joint uncertainty in X and Y

$$H(Y|X) + H(X) = H(X, Y)$$

$$H(X|Y) + \underline{H(Y)} = \underline{H(X, Y)}$$

Example:

$$P_{ij} = P(X = s_i, Y = r_j)$$

		$\frac{1}{2}$ r_0	$\frac{1}{4}$ r_1	$\frac{1}{8}$ r_2	$\frac{1}{8}$ r_3	
$\frac{1}{4}$ s_0		$\frac{1}{8}$	$\frac{1}{16}$	$\frac{1}{32}$	$\frac{1}{32}$	P_{03} $= P(X = s_0, Y = r_3)$ $= \frac{1}{32}$
$\frac{1}{4}$ s_1		$\frac{1}{16}$	$\frac{1}{8}$	$\frac{1}{32}$	$\frac{1}{32}$	
$\frac{1}{4}$ s_2		$\frac{1}{16}$	$\frac{1}{16}$	$\frac{1}{16}$	$\frac{1}{16}$	
$\frac{1}{4}$ s_3		$\frac{1}{4}$	0	0	0	

$P(Y|X)$.

$$\begin{aligned}
 & P_r(Y=r_1 | X=s_0) \\
 &= \frac{P_r(s_0, r_1)}{P_r(s_0)} \\
 &= \frac{\frac{1}{16}}{\frac{1}{4}} \\
 &= \frac{1}{4}
 \end{aligned}$$

$$P(Y|X=s_0).$$

$$P(Y=r_j|X=s_0).$$

$$= \frac{P_r(Y=r_j, X=s_0)}{P_r(X=s_0)}$$

$$P_r(Y=r_0|X=s_0) = \frac{\frac{1}{8}}{\frac{1}{4}} = \frac{1}{2}$$

$$P_r(Y=r_1|X=s_0) = \frac{\frac{1}{16}}{\frac{1}{4}} = \frac{1}{4}$$

$$P_r(Y=r_2|X=s_0) = \frac{1}{8}$$

$$= P_r(Y=r_3|X=s_0).$$

$$\begin{aligned}
 H(Y|X=s_0) &= H\left(\frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \frac{1}{8}\right) \\
 &= \frac{1}{2} \cdot \log_2 2 + \frac{1}{4} \cdot \log_2 4 \\
 &\quad + \left(\frac{1}{8} \cdot \log_2 8\right) \times 2 \\
 &= 1.75 \text{ bits}
 \end{aligned}$$

$$\begin{aligned}
 H(Y|X=s_1) &= H\left(\frac{1}{4}, \frac{1}{2}, \frac{1}{8}, 0\right) \\
 &= 1.75
 \end{aligned}$$

$$\begin{aligned}
 H(Y|X=s_2) &= H\left(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}\right) \\
 &= \frac{1}{4}(\log_2 4) \times 4 \\
 &= 2
 \end{aligned}$$

$$\begin{aligned}
 H(Y|X=s_3) &= H(1, 0, 0, 0) \\
 &= 0
 \end{aligned}$$

$$H(Y|X) = \sum_{i=1}^{M-1} \frac{Pr(X=s_i)}{H(Y|X=s_i)}$$

$$= \frac{1}{4} \times 1.75 + \frac{1}{4} \times 1.75 + \frac{1}{4} \times 2 + \frac{1}{4} \times 0$$

$$= \frac{22}{16} = \frac{11}{8} = 1.375 \text{ bits}$$

$H(Y|X)$

$$H(X) = 2 \text{ bits} = H\left(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}\right)$$

$$H(X) + H(Y|X) = 2 + 1.375 = 3.375$$

Joint Entropy of X, Y

$$= H(X, Y)$$

$$H(Y) + H(X|Y) = H(X, Y)$$

check

$$H(X) + H(Y|X) = H(X, Y)$$

information
in X

Conditional
Entropy

information in
 Y given X

Joint
information
in X, Y .