

Module 10: Finite Difference Methods for Boundary Value Problems

Lecture 39: Analytic Expression of the Error

The Lecture Contains:

 [More general boundary conditions](#)

 **Previous** **Next** 

Module 10: Finite Difference Methods for Boundary Value Problems

Lecture 39: Analytic Expression of the Error

Theorem: Let $p(x)$ and $q(x)$ satisfy the said inequalities and h satisfy $h \leq \frac{2}{p^*}$.

In addition, let $p(x), q(x)$ and $r(x)$ be so smooth that $y(x)$, the solution of the given BVP has a continuous sixth derivative on $[a, b]$. Then (10.8) holds with $\{u_j\}$, the solution of the difference system, and $e(x)$, the solution of the BVP

$$L_h e(x) = \theta(x), \quad a < x < b \quad (10.9)$$

$$e(a) = e(b) = 0$$

where

$$\theta(x) \equiv -\frac{1}{12} [y^{iv}(x) - 2p(x)y'''(x)].$$

Proof: We first note that the BVP (10.9) has a unique solution. Now define the net function $\{e_j\}$ by

$$e_j \equiv h^{-2} [y(x_j) - u_j] \quad 0 \leq j \leq J+1$$

using the given BVP, difference system, and the T.E. we obtain

$$\begin{aligned} L_h e_j &= h^{-2} [L_h y(x_j) - L_h u_j] \\ &= h^{-2} [L_h y(x_j) - L y(x_j)] + h^{-2} [L y(x_j) - L_h u_j] \\ &= h^{-2} T_j[y] \quad 1 \leq j \leq J. \end{aligned}$$

Now the T.E. can be written as

$$T_j[y] = h^2 \theta(x_j) + O(h^4) \text{ (using the continuity of } y^{vi}(x)).$$

Thus we have

$$L_h e_j = \theta(x_j) + O(h^2) \quad 1 \leq j \leq J$$

◀ Previous Next ▶

Module 10: Finite Difference Methods for Boundary Value Problems

Lecture 39: Analytic Expression of the Error

In addition

$$e_0 = e_{J+1} = 0 \quad \left[\begin{array}{l} \because y(a) = \alpha, y(b) = \beta \\ u_0 = \alpha, u_{J+1} = \beta \end{array} \right]$$

$$\text{Now } L_h[e(x_j) - e_j] = L_h e(x_j) - L e(x_j) - O(h^2)$$

$$= T_j[e] + O(h^2)$$

$$= O(h^2) \quad 1 \leq j \leq J$$

where we have used the smoothness of $e(x)$ in estimating $T(e)$.

From the stability of L_h , we get

$$|e(x_j) - e_j| \leq O(h^2) \quad (10.10)$$

Then multiplying by h^2 yields (10.8).

◀ Previous Next ▶

Module 10: Finite Difference Methods for Boundary Value Problems

Lecture 39: Analytic Expression of the Error

More general boundary conditions

Let us consider the BVP

$$L y(x) \equiv -y'' + p(x)y' + q(x)y = r(x) \quad (10.11)$$

subject to

$$\begin{aligned} a_0 y(a) - a_1 y'(a) &= \alpha & a_0 \geq 0, a_1 \geq 0 \\ b_0 y(b) + b_1 y'(b) &= \beta & b_0 \geq 0, b_1 \geq 0 & \quad a_0 + b_0 \neq 0 \end{aligned} \quad (10.12)$$

Let the difference equation, be

$$L_h u_j = r(x_j) \quad 0 \leq j \leq J+1$$

$$a_0 u_0 - a_1 \frac{u_1 - u_{-1}}{2h} = \alpha$$

$$b_0 u_{J+1} + b_1 \frac{u_{J+2} - u_J}{2h} = \beta$$

Again, we can show that for $h < \frac{2}{P^*}$, the difference system has a unique solution.

◀ Previous Next ▶