

MODULE 6

RANDOM VECTOR AND ITS JOINT DISTRIBUTION

LECTURES 25-36

Topics

6.1 MULTIVARIATE DISTRIBUTIONS

6.2 TYPES OF RANDOM VECTORS

6.3 CONDITIONAL DISTRIBUTIONS

6.4 INDEPENDENT RANDOM VARIABLES

6.5 EXPECTATIONS AND MOMENTS

6.5.1 Cauchy- Schwarz Inequality Random Variables

6.6 JOINT MOMENT GENERATING FUNCTION

6.7 PROPERTIES OF RANDOM VECTORS HAVING THE SAME DISTRIBUTION

6.7.1 Uniqueness Theorem

6.8 MULTINOMIAL DISTRIBUTION

6.8.1 Multinomial Distribution

6.9 BIVARIATE NORMAL DISTRIBUTION

6.10 DISTRIBUTION OF FUNCTION OF RANDOM VECTORS

6.10.1 Distribution Function Technique

6.10.1.1 Marginal Distribution of Order Statistics of a Random Sample Of Absolutely Continuous Type Random Variables

6.10.2 Transformation of Variables Technique

6.10.2.1 Distribution of Order Statistics

6.10.2.2 Distribution of Normalized Spacings of Exponential Distribution

6.10.3 Moment Generating Function Technique

6.11 DISTRIBUTIONS BASED ON SAMPLING FROM A NORMAL DISTRIBUTION

MODULE 6**RANDOM VECTOR AND ITS JOINT DISTRIBUTION****LECTURE 25****Topics****6.1 MULTIVARIATE DISTRIBUTIONS****6.1 MULTIVARIATE DISTRIBUTIONS**

A (univariate) random variable describes a numerical characteristic of a typical outcome of a random experiment. In many situations we may be interested in simultaneously studying two or more numerical characteristics of outcomes of a random experiment. To make the above discussion more clear consider the following example.

Example 1.1

Two distinguishable dice (labeled as D_1 and D_2) are thrown simultaneously. Here the sample space is $\Omega = \{(i, j) : i, j \in \{1, \dots, 6\}\}$, where the outcome $(i, j) \in \Omega$ indicates that i number of dots are observed on the uppermost face of die D_1 and j number of dots are observed on uppermost face of die D_2 . For $(i, j) \in \Omega$, define

$$X_1((i, j)) = i + j = \text{sum of number of dots on uppermost faces of two dice}$$

and

$$X_2((i, j)) = |i - j| = \text{absolute difference of number of dots on uppermost faces of two dice.}$$

It may be of interest to study numerical characteristics X_1 and X_2 simultaneously. This amounts to the study of the function $\underline{X} = (X_1, X_2) : \Omega \rightarrow \mathbb{R}$ defined on the sample space Ω . ■

Throughout $\mathbb{R}^p = \{\underline{x} = (x_1, \dots, x_p) : -\infty < x_i < \infty, i = 1, \dots, p\}$ will denote the p -dimensional Euclidean space and, for a set $B \subseteq \mathbb{R}^p$ and a function $\underline{X} = (X_1, X_2, \dots, X_p) : \Omega \rightarrow \mathbb{R}^p$,

$$\underline{X}^{-1}(B) \stackrel{\text{def}}{=} \{\omega \in \Omega : \underline{X}(\omega) = (X_1(\omega), X_2(\omega), \dots, X_p(\omega)) \in B\}.$$

Let (Ω, \mathcal{F}, P) be a given probability space.

Definition 1.1

A function $\underline{X} = (X_1, \dots, X_p): \Omega \rightarrow \mathbb{R}^p$ is called a p -dimensional *random vector* (or simple a random vector) if $\underline{X}^{-1}((-\infty, \underline{a}]) \in \mathcal{F}, \forall \underline{a} = (a_1, \dots, a_p) \in \mathbb{R}^p$; here $(-\infty, \underline{a}] = (-\infty, a_1] \times \dots \times (-\infty, a_p]$. ■

A 1-dimensional random vector will simply be referred to as a random variable. Clearly, a function $\underline{X} = (X_1, \dots, X_p): \Omega \rightarrow \mathbb{R}^p$ is a random vector if

$$\{\omega \in \Omega : X_1(\omega) \leq a_1, \dots, X_p(\omega) \leq a_p\} \in \mathcal{F}, \quad \forall \underline{a} = (a_1, \dots, a_p) \in \mathbb{R}^p.$$

For $\underline{a} = (a_1, \dots, a_p) \in \mathbb{R}^p$, $\underline{b} = (b_1, \dots, b_p) \in \mathbb{R}^p$, and $a_i < b_i, i = 1, \dots, p$, define

$$(\underline{a}, \underline{b}) = (a_1, b_1) \times \dots \times (a_p, b_p) \equiv \prod_{i=1}^p (a_i, b_i),$$

$$(\underline{a}, \underline{b}] = (a_1, b_1] \times \dots \times (a_p, b_p] \equiv \prod_{i=1}^p (a_i, b_i],$$

$$[\underline{a}, \underline{b}) = [a_1, b_1) \times \dots \times [a_p, b_p) \equiv \prod_{i=1}^p [a_i, b_i),$$

$$[\underline{a}, \underline{b}] = [a_1, b_1] \times \dots \times [a_p, b_p] \equiv \prod_{i=1}^p [a_i, b_i],$$

$$(-\infty, \underline{b}) = (-\infty, b_1) \times \dots \times (-\infty, b_p) \equiv \prod_{i=1}^p (-\infty, b_i),$$

$$(\underline{a}, \infty) = (a_1, \infty) \times \dots \times (a_p, \infty) \equiv \prod_{i=1}^p (a_i, \infty),$$

and

$$[\underline{a}, \infty) = [a_1, \infty) \times \dots \times [a_p, \infty) \equiv \prod_{i=1}^p [a_i, \infty).$$

Further define

$$\mathcal{C}_0 = \{(-\infty, \underline{b}] : \underline{b} \in \mathbb{R}^p\},$$

$$\mathcal{C}_1 = \{(\underline{a}, \underline{b}) : \underline{a}, \underline{b} \in \mathbb{R}^p, a_i < b_i, i = 1, \dots, p\},$$

$$\mathcal{C}_2 = \{(\underline{a}, \underline{b}] : \underline{a}, \underline{b} \in \mathbb{R}^p, a_i < b_i, i = 1, \dots, p\},$$

$$\mathcal{C}_3 = \{[\underline{a}, \underline{b}) : \underline{a}, \underline{b} \in \mathbb{R}^p, a_i < b_i, i = 1, \dots, p\},$$

$$\mathcal{C}_4 = \{[\underline{a}, \underline{b}] : \underline{a}, \underline{b} \in \mathbb{R}^p, a_i < b_i, i = 1, \dots, p\},$$

$$\mathcal{C}_5 = \{(-\infty, \underline{b}) : \underline{b} \in \mathbb{R}^p\},$$

$$\mathcal{C}_6 = \{(\underline{a}, \infty) : \underline{a} \in \mathbb{R}^p\},$$

and

$$\mathcal{C}_7 = \{[\underline{a}, \infty) : \underline{a} \in \mathbb{R}^p\}.$$

As in the case of $p = 1$ it can be shown that

- (i) \mathcal{B}_p = the Borel σ -field in $\mathbb{R}^p = \sigma(\mathcal{C}_i), i = 0, 1, \dots, 7$;
- (ii) $\{\underline{a}\} \in \mathcal{B}_p, \forall \underline{a} \in \mathbb{R}^p$, i. e., \mathcal{B}_p contains all singleton subsets of \mathbb{R}^p ;
- (iii) If $B \subseteq \mathbb{R}^p$ is countable then $B \in \mathcal{B}_p$;
- (iv) There exists a set $A \subseteq \mathbb{R}^p$ such that $A \notin \mathcal{B}_p$;
- (v) $\underline{X}: \Omega \rightarrow \mathbb{R}^p$ is a p -dimensional random vector if, and only if one of the following equivalent conditions hold:
 - a) $\underline{X}^{-1}(B) \in \mathcal{F}, \forall B \in \mathcal{C}_1$;
 - b) $\underline{X}^{-1}(B) \in \mathcal{F}, \forall B \in \mathcal{C}_2$;
 - c) $\underline{X}^{-1}(B) \in \mathcal{F}, \forall B \in \mathcal{C}_3$;
 - d) $\underline{X}^{-1}(B) \in \mathcal{F}, \forall B \in \mathcal{C}_4$;
 - e) $\underline{X}^{-1}(B) \in \mathcal{F}, \forall B \in \mathcal{C}_5$;
 - f) $\underline{X}^{-1}(B) \in \mathcal{F}, \forall B \in \mathcal{C}_6$;
 - g) $\underline{X}^{-1}(B) \in \mathcal{F}, \forall B \in \mathcal{C}_7$;
 - h) $\underline{X}^{-1}(B) \in \mathcal{F}, \forall B \in \mathcal{B}_p$.

- (vi) If $\underline{X} = (X_1, \dots, X_p)$ is a p -dimensional random vector and $g_i: \mathbb{R}^p \rightarrow \mathbb{R}, i = 1, \dots, k$, are Borel functions (i.e., $g_i^{-1}(B) \in \mathcal{F}, \forall B \in \mathcal{B}_p, i = 1, \dots, k$) then $(g_1(\underline{X}), \dots, g_k(\underline{X}))$ is a k -dimensional random vector.
- (vii) If $\underline{X}: \Omega \rightarrow \mathbb{R}^p$ is a p -dimensional random vector then $\underline{X}^{-1}(\{\underline{a}\}) = \{\omega \in \Omega: X_1(\omega) = a_1, \dots, X_p(\omega) = a_p\} \in \mathcal{F}, \forall \underline{a} = (a_1, \dots, a_p) \in \mathbb{R}^p$;
- (viii) The function $P_{\underline{X}}: \mathcal{B}_p \rightarrow \mathbb{R}$ given by,

$$P_{\underline{X}}(B) = P(\underline{X}^{-1}(B)), B \in \mathcal{B}_p,$$

is a probability measure on \mathcal{B}_p (i.e., $(\mathbb{R}^p, \mathcal{B}_p, P_{\underline{X}})$ is a probability space), called the *probability measure induced by \underline{X}* .

Example 1.2

Let $A, B \subseteq \Omega$. Define $\underline{X} = (X_1, X_2): \Omega \rightarrow \mathbb{R}^2$ by

$$X_1(\omega) = I_A(\omega) = \begin{cases} 1, & \text{if } \omega \in A \\ 0, & \text{if } \omega \notin A \end{cases};$$

and

$$X_2(\omega) = I_B(\omega) = \begin{cases} 1, & \text{if } \omega \in B \\ 0, & \text{if } \omega \notin B \end{cases}.$$

Then, for $\underline{a} = (a_1, a_2) \in \mathbb{R}^2$,

$$\begin{aligned} \underline{X}^{-1}((-\infty, \underline{a}]) &= \{\omega \in \Omega: X_1(\omega) \leq a_1, X_2(\omega) \leq a_2\} \\ &= \begin{cases} \phi, & \text{if } a_1 < 0 \text{ or } a_2 < 0 \\ A^c \cap B^c, & \text{if } 0 \leq a_1 < 1, 0 \leq a_2 < 1 \\ A^c, & \text{if } 0 \leq a_1 < 1, a_2 \geq 1 \\ B^c, & \text{if } a_1 \geq 1, 0 \leq a_2 < 1 \\ \Omega, & \text{if } a_1 \geq 1, a_2 \geq 1 \end{cases}. \end{aligned}$$

Thus

$$\begin{aligned} \underline{X} \text{ is a random vector} &\Leftrightarrow \underline{X}^{-1}((-\infty, \underline{a}]) \in \mathcal{F}, \forall \underline{a} \in \mathbb{R}^2 \\ &\Leftrightarrow A^c, B^c \in \mathcal{F} \\ &\Leftrightarrow A, B \in \mathcal{F} \end{aligned}$$

Thus \underline{X} is a random vector if, and only if, $A, B \in \mathcal{F}$. ■

Theorem 1.1

Let $\underline{X} = (X_1, \dots, X_p): \Omega \rightarrow \mathbb{R}^p$ be a given function. Then \underline{X} is a random vector if, and only if X_1, \dots, X_p ($X_i: \Omega \rightarrow \mathbb{R}, i = 1, \dots, p$) are random variables.

Proof. First suppose that $\underline{X} = (X_1, \dots, X_p)$ is a random vector. Then, for $a \in \mathbb{R}$, and for fixed $i \in \{1, \dots, p\}$

$$\begin{aligned} X_i^{-1}((-\infty, a]) &= \bigcap_{n=1}^{\infty} \underbrace{X_i^{-1}((-\infty, n] \times \dots \times (-\infty, n] \times (-\infty, a] \times (-\infty, n] \times \dots \times (-\infty, n]))}_{\in \mathcal{F}, \forall n=1,2,\dots}, \\ &\in \mathcal{F} \end{aligned}$$

i.e., X_i is a random variable.

Conversely suppose that X_1, \dots, X_p are random variables. Then, for $\underline{a} = (a_1, \dots, a_p) \in \mathbb{R}^p$,

$$\begin{aligned} \underline{X}^{-1}((-\infty, \underline{a}]) &= \{\omega \in \Omega: X_i(\omega) \leq a_i, i = 1, \dots, p\} \\ &= \bigcap_{i=1}^p \{\omega \in \Omega: X_i(\omega) \leq a_i\} \\ &= \bigcap_{i=1}^p \underbrace{X_i^{-1}((-\infty, a_i])}_{\in \mathcal{F}} \\ &\in \mathcal{F} \end{aligned}$$

i.e., \underline{X} is a random vector. ■

Remark 1.1

When Ω is countable we have $\mathcal{F} = \mathcal{P}(\Omega)$ and, therefore, any function $\underline{X} = (X_1, \dots, X_p): \Omega \rightarrow \mathbb{R}^p$ is a random vector. ■

Definition 1.2

- (i) The *joint distribution function* of a p -dimensional random vector $\underline{X} = (X_1, \dots, X_p): \Omega \rightarrow \mathbb{R}^p$ is defined by

$$F_{\underline{X}}(x_1, \dots, x_p) = P(\{\omega \in \Omega: X_1(\omega) \leq x_1, \dots, X_p(\omega) \leq x_p\}), \quad \underline{x} = (x_1, \dots, x_p) \in \mathbb{R}^p.$$

- (ii) The joint distribution function of any subset of random variables X_1, \dots, X_p is called a *marginal distribution function* of $F_{\underline{X}}(\cdot)$. ■

Remark 1.2

- (i) If $F_{\underline{X}}(\cdot)$ is the distribution function of a p -dimensional random vector $\underline{X} = (X_1, \dots, X_p)$ then

$$\begin{aligned}
 F_{\underline{X}}(\underline{x}) &= P(\{X_i \leq x_i, i = 1, \dots, p\}) \\
 &= P\left(\underline{X}^{-1}\left((-\infty, \underline{x}]\right)\right) \\
 &= P_{\underline{X}}\left((-\infty, \underline{x}]\right) \\
 &= P\left(\bigcap_{i=1}^p \{X_i \leq x_i\}\right) \\
 &= P\left(\bigcap_{i=1}^p X_i^{-1}\left((-\infty, x_i]\right)\right), \quad \underline{x} = (x_1, \dots, x_p) \in \mathbb{R}^p.
 \end{aligned}$$

- (ii) Let $F_{X_1, \dots, X_p}(\cdot)$ be the distribution function of a random vector $\underline{X} = (X_1, \dots, X_p)$ and let $\underline{\beta} = (\beta_1, \dots, \beta_p)$ be a permutation of $(1, \dots, p)$. Then

$$\begin{aligned}
 F_{X_1, \dots, X_p}(x_1, \dots, x_p) &= P\left(\bigcap_{i=1}^p \{X_i \leq x_i\}\right) \\
 &= P\left(\bigcap_{i=1}^p \{X_{\beta_i} \leq x_{\beta_i}\}\right) \\
 &= F_{X_{\beta_1}, \dots, X_{\beta_p}}(x_{\beta_1}, \dots, x_{\beta_p}), \quad \underline{x} = (x_1, \dots, x_p) \in \mathbb{R}^p.
 \end{aligned}$$

It follows that the distribution function of $(X_{\beta_1}, \dots, X_{\beta_p})$ is given by

$F_{X_{\beta_1}, \dots, X_{\beta_p}}(y_1, \dots, y_p) = F_{X_1, \dots, X_p}(y_{\gamma_1}, \dots, y_{\gamma_p})$, $\underline{y} = (y_1, \dots, y_p) \in \mathbb{R}^p$, where $\underline{\gamma} = (\gamma_1, \dots, \gamma_p)$ is the inverse permutation of $\underline{\beta} = (\beta_1, \dots, \beta_p)$. To illustrate this point, consider $p = 3$ and $\underline{\beta} = (\beta_1, \beta_2, \beta_3) = (2, 3, 1)$. Then the

inverse permutation of $\underline{\beta} = (\beta_1, \beta_2, \beta_3)$ is $\underline{\gamma} = (\gamma_1, \gamma_2, \gamma_3) = (3, 1, 2)$, and therefore, for $\underline{y} = (y_1, y_2, y_3) \in \mathbb{R}^3$,

$$\begin{aligned}
 F_{X_{\beta_1}, X_{\beta_2}, X_{\beta_3}}(y_1, y_2, y_3) &= F_{X_2, X_3, X_1}(y_1, y_2, y_3) \\
 &= P(\{X_2 \leq y_1, X_3 \leq y_2, X_1 \leq y_3\}) \\
 &= P(\{X_1 \leq y_3, X_2 \leq y_1, X_3 \leq y_2\}) \\
 &= F_{X_1, X_2, X_3}(y_3, y_1, y_2) \\
 &= F_{X_1, X_2, X_3}(\gamma_1, \gamma_2, \gamma_3).
 \end{aligned}$$

(iii) Note that a distribution function $F_{X_1, \dots, X_p}(x_1, \dots, x_p)$ is increasing in each argument when other arguments are kept fixed.

We recall the following results from the theory of multivariable calculus.

Lemma 1.1

Let $D \subseteq \mathbb{R}^p$ and let $g: D \rightarrow \mathbb{R}$ be a function such that:

- (i) g is bounded above, i.e., there exists a real constant M such that $g(\underline{x}) \leq M, \forall \underline{x} \in D$;
- (ii) for every fixed $i \in \{1, \dots, p\}$ and fixed $(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_p) \in \mathbb{R}^{p-1}$, $g(x_1, \dots, x_{i-1}, t, x_{i+1}, \dots, x_p)$ is non decreasing in $t \in D_i = \{y \in \mathbb{R}: (x_1, \dots, x_{i-1}, y, x_{i+1}, \dots, x_p) \in D\}$. Then $\lim_{\underline{x} \rightarrow \underline{\infty}} g(\underline{x})$ exists and, for any permutation $\underline{\beta} = (\beta_1, \dots, \beta_p)$ of $(1, \dots, p)$,

$$\lim_{x_{\beta_p} \rightarrow \infty} \cdots \lim_{x_{\beta_1} \rightarrow \infty} g(x_1, \dots, x_p) = \lim_{\underline{x} \rightarrow \underline{\infty}} g(\underline{x}).$$

In particular all iterated limits

$$\lim_{x_{\beta_p} \rightarrow \infty} \cdots \lim_{x_{\beta_1} \rightarrow \infty} g(x_1, \dots, x_p), \quad (\beta_1, \dots, \beta_p) \in S_p,$$

exist and are equal, where S_p denotes the set of all permutations of $(1, \dots, p)$. We denote the common value of all iterated limits by

$$\lim_{\substack{x_i \rightarrow \infty \\ i=1, \dots, p}} g(\underline{x}). \blacksquare$$

Note that if $F_{\underline{X}}(\cdot)$ is a distribution function in \mathbb{R}^p ($p \geq 2$) then, for a fixed $k \in \{1, \dots, p-1\}$ and fixed $(x_{k+1}, \dots, x_p) \in \mathbb{R}^{p-k}$, the function $g: \mathbb{R}^k \rightarrow \mathbb{R}$, given by

$$g(x_1, \dots, x_k) = F_{\underline{X}}(x_1, \dots, x_k, x_{k+1}, \dots, x_p),$$

satisfies properties (i) and (ii) stated in Lemma 1.1. Therefore, for fixed $(x_{k+1}, \dots, x_p) \in \mathbb{R}^{p-k}$

$$\lim_{\underline{x}^* \rightarrow \underline{\infty}} F_{\underline{X}}(x_1, \dots, x_k, x_{k+1}, \dots, x_p) = \lim_{\substack{x_i \rightarrow \infty \\ i=1, \dots, k}} F_{\underline{X}}(x_1, \dots, x_k, x_{k+1}, \dots, x_p),$$

where $\underline{x}^* = (x_1, \dots, x_k)$.

Lemma 1.2

Let $F_{\underline{X}}(\cdot)$ be the distribution function of a p -dimensional ($p \geq 2$) random vector $\underline{X} = (X_1, \dots, X_p)$. For a fixed positive integer $k \in \{1, \dots, p-1\}$, let $\underline{Y} = (X_1, \dots, X_k)$ and let $\underline{Z} = (X_{k+1}, \dots, X_p)$ so that $\underline{X} = (\underline{Y}, \underline{Z})$. Then the marginal distribution function of $\underline{Y} = (Y_1, \dots, Y_k)$ is given by

$$F_{\underline{Y}}(x_1, \dots, x_k) = \lim_{\substack{x_i \rightarrow \infty \\ i=k+1, \dots, p}} F_{\underline{X}}(x_1, \dots, x_k, x_{k+1}, \dots, x_p), \quad (x_1, \dots, x_k) \in \mathbb{R}^k.$$

Proof. For fixed $x_1, \dots, x_{p-1} \in \mathbb{R}$

$$\begin{aligned} \lim_{x_p \rightarrow \infty} F_{\underline{X}}(x_1, \dots, x_p) &= \lim_{x_p \rightarrow \infty} P \left(\bigcap_{i=1}^p X_i^{-1}((-\infty, x_i]) \right) \\ &= \lim_{n \rightarrow \infty} P \left(\underbrace{\left(\bigcap_{i=1}^{p-1} X_i^{-1}((-\infty, x_i]) \right)}_{=A_n \uparrow} \bigcap X_p^{-1}((-\infty, n]) \right) \\ &= P \left(\bigcup_{n=1}^{\infty} A_n \right) \\ &= P \left(\bigcap_{i=1}^{p-1} X_i^{-1}((-\infty, x_i]) \right) \quad \text{since } \left(\bigcup_{n=1}^{\infty} X_p^{-1}((-\infty, n]) \right) = \Omega \end{aligned}$$

$$= F_{X_1, \dots, X_{p-1}}(x_1, \dots, x_{p-1}). \quad (1.1)$$

Now the assertion follows on recursively using (1.1). ■

Remark 1.3

Let $\underline{X} = (X_1, \dots, X_p)$ be a random vector and let $\underline{\beta} = (\beta_1, \dots, \beta_p) \in S_p$, the set of all permutations of $(1, \dots, p)$. If $\underline{\gamma} = (\gamma_1, \dots, \gamma_p)$ is the inverse permutation of $(\beta_1, \dots, \beta_p)$ then, for a fixed $k \in \{1, \dots, p-1\}$, the marginal distribution function of $(X_{\beta_1}, \dots, X_{\beta_k})$ is given by

$$\begin{aligned} F_{X_{\beta_1}, \dots, X_{\beta_k}}(x_1, \dots, x_k) &= \lim_{\substack{x_j \rightarrow \infty \\ j=k+1, \dots, p}} F_{X_{\beta_1}, \dots, X_{\beta_p}}(x_1, \dots, x_p) \\ &= \lim_{\substack{x_j \rightarrow \infty \\ j=k+1, \dots, p}} F_{X_1, \dots, X_p}(x_{\gamma_1}, \dots, x_{\gamma_p}). \blacksquare \end{aligned}$$

Let $\underline{X} = (X_1, \dots, X_p)$ be a random vector and let $\underline{a} = (a_1, \dots, a_p)$, $\underline{b} = (b_1, \dots, b_p) \in \mathbb{R}^p$. Then

$$\begin{aligned} P(\{a_1 < X_1 \leq b_1\}) &= P(\{X_1 \leq b_1\}) - P(\{X_1 \leq a_1\}) \\ &= F_{X_1}(b_1) - F_{X_1}(a_1). \end{aligned} \quad (1.2)$$

Also

$$\begin{aligned} P(\{a_1 < X_1 \leq b_1, a_2 < X_2 \leq b_2\}) &= P(\{a_1 < X_1 \leq b_1, X_2 \leq b_2\}) \\ &\quad - P(\{a_1 < X_1 \leq b_1, X_2 \leq a_2\}) \\ &= [P(\{X_1 \leq b_1, X_2 \leq b_2\}) - P(\{X_1 \leq a_1, X_2 \leq b_2\})] \\ &\quad - [P(\{X_1 \leq b_1, X_2 \leq a_2\}) - P(\{X_1 \leq a_1, X_2 \leq a_2\})] \\ &= F_{X_1, X_2}(b_1, b_2) - [F_{X_1, X_2}(a_1, b_2) + F_{X_1, X_2}(b_1, a_2)] \\ &\quad + F_{X_1, X_2}(a_1, a_2). \end{aligned} \quad (1.3)$$

To write the expression of $P(\{a_i < X_i \leq b_i, i = 1, \dots, p\})$ in a closed form define, for $k \in \{0, 1, \dots, p\}$,

$$\Delta_{k,p} \equiv \Delta_{k,p}(\underline{a}, \underline{b}) = \{\underline{z} \in \mathbb{R}^p : z_i \in \{a_i, b_i\}, i = 1, \dots, p, \text{ and } k \text{ of } z_1, \dots, z_p \text{ are } a_j\text{'s}\}. \quad (1.4)$$

Note that the set $\Delta_{k,p}$ has $\binom{p}{k}$ elements. From (1.2) and (1.3) we have

$$P(\{a_1 < X_1 \leq b_1\}) = F_{X_1}(b_1) - F_{X_1}(a_1) = \sum_{k=0}^1 (-1)^k \sum_{z \in \Delta_{k,1}} F_{X_1}(z) \quad (1.5)$$

and

$$P(\{a_i < X_i \leq b_i, i = 1, 2\}) = \sum_{k=0}^2 (-1)^k \sum_{(z_1, z_2) \in \Delta_{k,2}} F_{X_1, X_2}(z_1, z_2) \quad (1.6)$$

Lemma 1.3

Let $\underline{X} = (X_1, \dots, X_p): \Omega \rightarrow \mathbb{R}^p$ be a random vector and let $\underline{a} = (a_1, \dots, a_p)$, $\underline{b} = (b_1, \dots, b_p) \in \mathbb{R}^p$. Let $\Delta_{k,p} \equiv \Delta_{k,p}((\underline{a}, \underline{b}])$, $k = 0, 1, \dots, p$ be as defined in (1.4). Then

$$P(\{a_i < X_i \leq b_i, i = 1, \dots, p\}) = \sum_{k=0}^p (-1)^k \sum_{\underline{z} \in \Delta_{k,p}((\underline{a}, \underline{b}])} F_{\underline{X}}(\underline{z}). \quad (1.7)$$

Proof. From (1.5) and (1.6) it is clear that the result is true for $p = 1$ and $p = 2$. Now suppose that (1.7) holds for general p -dimensional random vectors. For simplicity assume that $P(\{a_{p+1} < X_{p+1} \leq b_{p+1}\}) > 0$. Then, for $(X_1, \dots, X_p, X_{p+1}): \Omega \rightarrow \mathbb{R}^{p+1}$, $\underline{a} = (a_1, \dots, a_p) \in \mathbb{R}^p$, $\underline{b} = (b_1, \dots, b_p) \in \mathbb{R}^p$, $\underline{a}^* = (a_1, \dots, a_p, a_{p+1}) \in \mathbb{R}^{p+1}$ and $\underline{b}^* = (b_1, \dots, b_p, b_{p+1}) \in \mathbb{R}^{p+1}$.

$$\begin{aligned} & P(\{a_i < X_i \leq b_i, i = 1, \dots, p+1\}) \\ &= P(\{a_i < X_i \leq b_i, i = 1, \dots, p\} | \{a_{p+1} < X_{p+1} \leq b_{p+1}\}) P(\{a_{p+1} < X_{p+1} \leq b_{p+1}\}) \\ &= \sum_{k=0}^p (-1)^k \sum_{\underline{z} \in \Delta_{k,p}((\underline{a}, \underline{b}])} P(\{X_i \leq z_i, i = 1, \dots, p\} | \{a_{p+1} < X_{p+1} \leq b_{p+1}\}) P(\{a_{p+1} < X_{p+1} \leq b_{p+1}\}) \\ &= \sum_{k=0}^p (-1)^k \sum_{\underline{z} \in \Delta_{k,p}((\underline{a}, \underline{b}])} P(\{X_i \leq z_i, i = 1, \dots, p, a_{p+1} < X_{p+1} \leq b_{p+1}\}) \\ &= \sum_{k=0}^p (-1)^k \sum_{\underline{z} \in \Delta_{k,p}((\underline{a}, \underline{b}])} [P(\{X_1 \leq z_1, \dots, X_p \leq z_p, X_{p+1} \leq b_{p+1}\}) \\ &\quad - P(\{X_1 \leq z_1, \dots, X_p \leq z_p, X_{p+1} \leq a_{p+1}\})]. \end{aligned}$$

It is easy to verify that

$$\begin{aligned}
& \sum_{k=0}^p (-1)^k \sum_{\underline{z} \in \Delta_{k,p}(\underline{a}, \underline{b})} [P(\{X_1 \leq z_1, \dots, X_p \leq z_p, X_{p+1} \leq b_{p+1}\}) \\
& \quad - P(\{X_1 \leq z_1, \dots, X_p \leq z_p, X_{p+1} \leq a_{p+1}\})] \\
& = \sum_{k=0}^{p+1} (-1)^k \sum_{\underline{t} \in \Delta_{k,p+1}(\underline{a}^*, \underline{b}^*)} F_{X_1, \dots, X_{p+1}}(t_1, \dots, t_{p+1}),
\end{aligned}$$

and therefore the assertion follows by principle of mathematical induction. ■

Theorem 1.2

Let $F_{\underline{X}}(\cdot)$ be the distribution of a p -dimensional random vector $\underline{X} = (X_1, \dots, X_p)$. Then

- (i) $\lim_{\substack{x_i \rightarrow \infty \\ i=1, \dots, p}} F_{\underline{X}}(x_1, \dots, x_p) = 1;$
- (ii) For each fixed $i \in \{1, \dots, p\}$ and fixed $(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_p) \in \mathbb{R}^{p-1},$

$$\lim_{y \rightarrow -\infty} F_{\underline{X}}(x_1, \dots, x_{i-1}, y, x_{i+1}, \dots, x_p) = 0;$$

- (iii) $F_{\underline{X}}(x_1, \dots, x_p)$ is right continuous in each argument (keeping other arguments fixed);
- (iv) For each rectangle $(\underline{a}, \underline{b}] \in \mathbb{R}^p$

$$\sum_{k=0}^p (-1)^k \sum_{\underline{z} \in \Delta_{k,p}(\underline{a}, \underline{b})} F_{\underline{X}}(\underline{z}) \geq 0.$$

Proof. Note that, for $(\underline{a}, \underline{b}] \in \mathbb{R}^p,$

$$\sum_{k=0}^p (-1)^k \sum_{\underline{z} \in \Delta_{k,p}(\underline{a}, \underline{b})} F_{\underline{X}}(\underline{z}) = P(\underline{X} \in (\underline{a}, \underline{b}]) \geq 0. \quad (\text{using Lemma 1.3})$$

This proves (iv).

For notational convenience we will provide the proofs of (i) - (iii) for only $p = 2$.

- (i) For fixed $x_1 \in \mathbb{R}$

$$\begin{aligned}
\lim_{x_2 \rightarrow \infty} F_{X_1, X_2}(x_1, x_2) &= \lim_{n \rightarrow \infty} F_{X_1, X_2}(x_1, n) \\
&= \lim_{n \rightarrow \infty} P(\underbrace{X_1^{-1}((-\infty, x_1]) \cap X_2^{-1}((-\infty, n])}_{=A_n \uparrow})
\end{aligned}$$

$$\begin{aligned}
&= P\left(\bigcup_{n=1}^{\infty} A_n\right) \\
&= P(X_1^{-1}(-\infty, x_1]).
\end{aligned}$$

Therefore,

$$\begin{aligned}
\lim_{x_1 \rightarrow \infty} \lim_{x_2 \rightarrow \infty} F_{X_1, X_2}(x_1, x_2) &= \lim_{x_1 \rightarrow \infty} P(X_1^{-1}(-\infty, x_1]) \\
&= \lim_{n \rightarrow \infty} P\left(\underbrace{X_1^{-1}((-\infty, n])}_{=B_n \uparrow}\right) \\
&= P\left(\bigcup_{n=1}^{\infty} B_n\right) \\
&= P(\Omega) \\
&= 1.
\end{aligned}$$

(ii) Fix $x_2 \in \mathbb{R}$. Then

$$\begin{aligned}
\lim_{x_1 \rightarrow -\infty} F_{X_1, X_2}(x_1, x_2) &= \lim_{n \rightarrow \infty} P\left(\underbrace{X_1^{-1}(-\infty, -n] \cap X_2^{-1}((-\infty, x_2])}_{=B_n \downarrow}\right) \\
&= P\left(\bigcap_{n=1}^{\infty} B_n\right) \\
&= P(\phi) \\
&= 0.
\end{aligned}$$

Similarly, for fixed $x_1 \in \mathbb{R}$

$$\lim_{x_2 \rightarrow -\infty} F_{X_1, X_2}(x_1, x_2) = 0.$$

(iii) Let $\underline{x} = (x_1, x_2) \in \mathbb{R}^2$. Then

$$\begin{aligned}
\lim_{h \downarrow 0} F_{X_1, X_2}(x_1 + h, x_2) &= \lim_{n \rightarrow \infty} F_{X_1, X_2}\left(x_1 + \frac{1}{n}, x_2\right) \\
&= \lim_{n \rightarrow \infty} P\left(\underbrace{X_1^{-1}\left((-\infty, x_1 + \frac{1}{n}]\right) \cap X_2^{-1}((-\infty, x_2])}_{=C_n \downarrow}\right)
\end{aligned}$$

$$\begin{aligned}
&= P\left(\bigcap_{n=1}^{\infty} C_n\right) \\
&= P\left(X_1^{-1}((-\infty, x_1]) \cap X_2^{-1}((-\infty, x_2])\right) \\
&= F_{X_1, X_2}(x_1, x_2),
\end{aligned}$$

i.e., for every fixed $x_2 \in \mathbb{R}$, $F_{X_1, X_2}(x_1, x_2)$ is right continuous in x_1 . Similarly, for every fixed $x_1 \in \mathbb{R}$, $F_{X_1, X_2}(x_1, x_2)$ is right continuous in x_2 . ■

Remark 1.4

- (i) Let $\Delta_p = \bigcup_{k=0}^p \Delta_{k,p}$. Then Δ_p is the set of 2^p vertices of the rectangle $(\underline{a}, \underline{b}] \in \mathbb{R}^p$. For example, for $p = 1$, $(\underline{a}, \underline{b}] = (a_1, b_1]$, $\Delta_1 = \{a_1, b_1\}$ and, for $p = 2$, $(\underline{a}, \underline{b}] = (a_1, b_1] \times (a_2, b_2]$, $\Delta_2 = \{(b_1, b_2), (b_1, a_2), (a_1, b_2), (a_1, a_2)\}$.




Figure 1.1




Figure 1.2

- (ii) Note that, for $p = 1$, the assertion (iv) of Theorem 1.2 reduces to $F_X(b) \geq F_X(a), \forall -\infty < a \leq b < \infty$ i.e., F_X is non-decreasing.