

MODULE 3
FUNCTION OF A RANDOM VARIABLE AND ITS
DISTRIBUTION
LECTURES 12-16

Topics

3.1 FUNCTION OF A RANDOM VARIABLE

3.2 PROBABILITY DISTRIBUTION OF A FUNCTION OF A RANDOM VARIABLE

3.3 EXPECTATION AND MOMENTS OF A RANDOM VARIABLE

3.4 PROPERTIES OF RANDOM VARIABLES HAVING THE SAME DISTRIBUTION

3.5 PROBABILITY AND MOMENT INEQUALITIES

3.5.1 Markov Inequality

3.5.2 Chebyshev Inequality

3.5.3 Jensen Inequality

3.5.4 AM-GM-HM inequality

3.6 DESCRIPTIVE MEASURES OF PROBABILITY DISTRIBUTIONS

3.6.1 Measures of Central Tendency

3.6.1.1 Mean

3.6.1.2 Median

3.6.1.3 Mode

3.6.2 Measures of Dispersion

3.6.2.1 Standard Deviation

3.6.2.2 Mean Deviation

3.6.2.3 Quartile Deviation

3.6.2.4 Coefficient of Variation

3.7 MEASURES OF SKEWNESS**3.8 MEASURES OF KURTOSIS****MODULE 3****FUNCTION OF A RANDOM VARIABLE AND ITS DISTRIBUTION****LECTURE 12****Topics****3.1 FUNCTION OF A RANDOM VARIABLE****3.2 PROBABILITY DISTRIBUTION OF A FUNCTION OF A RANDOM VARIABLE****3.1 FUNCTION OF A RANDOM VARIABLE**

Let (Ω, \mathcal{F}, P) be a probability space and let X be random variable defined on (Ω, \mathcal{F}, P) . Further let $h: \mathbb{R} \rightarrow \mathbb{R}$ be a given function and let $Z: \Omega \rightarrow \mathbb{R}$ be a function of random variable X , defined by $Z(\omega) = h(X(\omega)), \omega \in \Omega$. In many situations it may be of interest to study the probabilistic properties of Z , which is a function of random variable X . Since the variable Z takes values in \mathbb{R} , to study the probabilistic properties of Z , it is necessary that $Z^{-1}(B) \in \mathcal{F}, \forall B \in \mathcal{B}_1$, i.e., Z is a random variable. Throughout, for a positive integer k , \mathbb{R}^k will denote the k -dimensional Euclidean space and \mathcal{B}_k will denote the Borel sigma-field in \mathbb{R}^k .

Definition 1.1

Let k and m be positive integers. A function $h: \mathbb{R}^k \rightarrow \mathbb{R}^m$ is said to be a Borel function if $h^{-1}(B) \in \mathcal{B}_k, \forall B \in \mathcal{B}_m$. ■

The following lemma will be useful in deriving conditions on the function $h: \mathbb{R} \rightarrow \mathbb{R}$ so that $Z: \Omega \rightarrow \mathbb{R}$, defined by $Z(\omega) = h(X(\omega))$, $\omega \in \Omega$, is a random variable. Recall that, for a function $\Psi: D_1 \rightarrow D_2$ and $A \subseteq D_2$, $\Psi^{-1}(A) = \{\omega \in D_1: \Psi(\omega) \in A\}$.

Lemma 1.1

Let $X: \Omega \rightarrow \mathbb{R}$ and $h: \mathbb{R} \rightarrow \mathbb{R}$ be given functions. Define $Z: \Omega \rightarrow \mathbb{R}$ by $Z(\omega) = h(X(\omega))$, $\omega \in \Omega$. Then, for any $B \subseteq \mathbb{R}$,

$$Z^{-1}(B) = X^{-1}(h^{-1}(B)).$$

Proof. Fix $B \subseteq \mathbb{R}$. Note that $h^{-1}(B) = \{x \in \mathbb{R}: h(x) \in B\}$. Clearly

$$h(X(\omega)) \in B \Leftrightarrow X(\omega) \in h^{-1}(B).$$

Therefore,

$$\begin{aligned} Z^{-1}(B) &= \{\omega \in \Omega: Z(\omega) \in B\} \\ &= \{\omega \in \Omega: h(X(\omega)) \in B\} \\ &= \{\omega \in \Omega: X(\omega) \in h^{-1}(B)\} \\ &= X^{-1}(h^{-1}(B)). \blacksquare \end{aligned}$$

Theorem 1.1

Let X be a random variable defined on a probability space (Ω, \mathcal{F}, P) and let $h: \mathbb{R} \rightarrow \mathbb{R}$ be a Borel function. Then the function $Z: \Omega \rightarrow \mathbb{R}$, defined by $Z(\omega) = h(X(\omega))$, $\omega \in \Omega$, is a random variable.

Proof. Fix $B \in \mathcal{B}_1$. Since h is a Borel function, we have $h^{-1}(B) \in \mathcal{B}_1$. Now using the fact that X is a random variable it follows that

$$Z^{-1}(B) = X^{-1}(h^{-1}(B)) \in \mathcal{F}.$$

This proves the result. \blacksquare

Remark 1.1

- (i) Let $h: \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function. According to a standard result in calculus inverse image of any open interval (a, b) , $-\infty \leq a < b \leq \infty$, under continuous function h is a countable union of disjoint open intervals. Since \mathcal{B}_1 contains all open intervals and is closed under countable unions it follows that $h^{-1}((a, b)) \in \mathcal{B}_1$, whenever $-\infty \leq a < b \leq \infty$. Now on employing the

arguments similar to the one used in proving Theorem 1.1, Module 2 (also see Theorem 1.2, Module 2) we conclude that $h^{-1}(B) \in \mathcal{B}_1, \forall B \in \mathcal{B}_1$. It follows that any continuous function $h: \mathbb{R} \rightarrow \mathbb{R}$ is a Borel function and thus, in view of Theorem 1.1, any continuous function of a random variable is a random variable. In particular if X is a random variable then $X^2, |X|, \max(X, 0), \sin X$ and $\cos X$ are random variables.

- (ii) Let $h: \mathbb{R} \rightarrow \mathbb{R}$ be a strictly monotone function. Then, for $-\infty \leq a < b \leq \infty$, $h^{-1}(a, b)$ is a countable union of intervals and therefore $h^{-1}(a, b) \in \mathcal{B}_1$, i.e., h is a Borel function. It follows that if X is a random variable and if $h: \mathbb{R} \rightarrow \mathbb{R}$ is strictly monotone then $h(X)$ is a random variable. ■

A random variable X takes values in various Borel sets according to some probability law called the probability distribution of random variable X . Clearly the probability distribution of a random variable of absolutely continuous/discrete type is described by its distribution function (d.f.) and/or by its probability density function/probability mass function (p.d.f/p.m.f.). For a given Borel function $h: \mathbb{R} \rightarrow \mathbb{R}$, in the following section, we will derive probability distribution of $h(X)$ using the probability distribution of random variable X . ■

3.2 PROBABILITY DISTRIBUTION OF A FUNCTION OF A RANDOM VARIABLE

In our future discussions when we refer to a random variable, unless otherwise stated, it will be either of discrete type or of absolutely continuous type. The probability distribution of a discrete type random variable will be referred to as a discrete (probability) distribution and the probability distribution of a random variable of absolutely continuous type will be referred to as an absolutely continuous (probability) distribution.

The following theorem deals with discrete probability distributions.

Theorem 2.1

Let X be a random variable of discrete type with support S_X and p.m.f. $f_X(\cdot)$. Let $h: \mathbb{R} \rightarrow \mathbb{R}$ be a Borel function and let $Z: \Omega \rightarrow \mathbb{R}$ be defined by $Z(\omega) = h(X(\omega)), \omega \in \Omega$. Then Z is a random variable of discrete type with support $S_Z = \{h(x): x \in S_X\}$ and p.m.f.

$$f_Z(z) = \begin{cases} \sum_{x \in A_z} f_X(x), & \text{if } z \in S_Z \\ 0, & \text{otherwise} \end{cases}$$

$$= \begin{cases} P(\{X \in A_z\}), & \text{if } z \in S_Z \\ 0, & \text{otherwise} \end{cases}$$

where $A_z = \{x \in S_X : h(x) = z\}$.

Proof. Since h is a Borel function, using Theorem 1.1, it follows that Z is a random variable. Also X is of discrete implies that S_X is countable which further implies that S_Z is countable. Fix $z_0 \in S_Z$, so that $z_0 = h(x_0)$ for some $x_0 \in S_X$.

Then

$$\begin{aligned} \{X = x_0\} &= \{\omega \in \Omega : X(\omega) = x_0\} \subseteq \{\omega \in \Omega : h(X(\omega)) = h(x_0)\} \\ &= \{h(X) = h(x_0)\} \\ &= \{Z = z_0\}, \end{aligned}$$

and

$$\begin{aligned} \{X \in S_X\} &= \{\omega \in \Omega : X(\omega) \in S_X\} \subseteq \{\omega \in \Omega : h(X(\omega)) \in S_Z\} \\ &= \{h(X) \in S_Z\} \\ &= \{Z \in S_Z\}. \end{aligned}$$

Therefore,

$$P(\{Z = z_0\}) \geq P(\{X = x_0\}) > 0, \quad (\text{since } x_0 \in S_X),$$

$$\text{and } P(\{Z \in S_Z\}) \geq P(\{X \in S_X\}) = 1.$$

It follows that S_Z is countable, $P(\{Z = z\}) > 0, \forall z \in S_Z$ and $P(\{Z \in S_Z\}) = 1$, i.e., Z is a discrete type random variable with support S_Z .

Moreover, for $z \in S_Z$,

$$\begin{aligned} P(\{Z = z\}) &= P(\{\omega \in \Omega : h(X(\omega)) = z\}) \\ &= \sum_{x \in A_z} P(\{X = x\}) \\ &= \sum_{x \in A_z} f_X(x) \\ &= P(\{X \in A_z\}). \end{aligned}$$

Hence the result follows. ■

The following corollary is an immediate consequence of the above theorem.

Corollary 2.1

Under the notation and assumptions of Theorem 2.1, suppose that $h: \mathbb{R} \rightarrow \mathbb{R}$ is one-one with inverse function $h^{-1}: D \rightarrow \mathbb{R}$, where $D = \{h(x): x \in \mathbb{R}\}$. Then Z is a discrete type random variable with support $S_Z = \{h(x): x \in S_X\}$ and p.m.f.

$$\begin{aligned} f_Z(z) &= \begin{cases} f_X(h^{-1}(z)), & \text{if } z \in S_Z \\ 0, & \text{otherwise} \end{cases} \\ &= \begin{cases} P(\{X = h^{-1}(z)\}), & \text{if } z \in S_Z \\ 0, & \text{otherwise} \end{cases} \blacksquare \end{aligned}$$

Example 2.1

Let X be a random variable with p.m.f.

$$f_X(x) = \begin{cases} \frac{1}{7}, & \text{if } x \in \{-2, -1, 0, 1\} \\ \frac{3}{14}, & \text{if } x \in \{2, 3\} \\ 0, & \text{otherwise} \end{cases}.$$

Show that $Z = X^2$ is a random variable. Find its p.m.f. and distribution function.

Solution. Since $h(x) = x^2, x \in \mathbb{R}$, is a continuous function and X is a random variable, using Remark 1.1 (i) it follows that $Z = h(X) = X^2$ is a random variable. Clearly $S_X = \{-2, -1, 0, 1, 2, 3\}$ and $S_Z = \{0, 1, 4, 9\}$. Moreover,

$$P(\{Z = 0\}) = P(\{X^2 = 0\}) = P(\{X = 0\}) = \frac{1}{7},$$

$$P(\{Z = 1\}) = P(\{X^2 = 1\}) = P(X \in \{-1, 1\}) = \frac{1}{7} + \frac{1}{7} = \frac{2}{7},$$

$$P(\{Z = 4\}) = P(\{X^2 = 4\}) = P(X \in \{-2, 2\}) = \frac{1}{7} + \frac{3}{14} = \frac{5}{14},$$

$$\text{and } P(\{Z = 9\}) = P(\{X^2 = 9\}) = P(\{X \in \{-3, 3\}\}) = 0 + \frac{3}{14} = \frac{3}{14}.$$

Therefore the p.m.f. of Z is

$$f_Z(z) = \begin{cases} \frac{1}{7}, & \text{if } z = 0 \\ \frac{2}{7}, & \text{if } z = 1 \\ \frac{5}{14}, & \text{if } z = 4 \\ \frac{3}{14}, & \text{if } z = 9 \\ 0, & \text{otherwise} \end{cases},$$

and the distribution function of Z is

$$F_Z(z) = \begin{cases} 0, & \text{if } z < 0 \\ \frac{1}{7}, & \text{if } 0 \leq z < 1 \\ \frac{3}{7}, & \text{if } 1 \leq z < 4 \\ \frac{11}{14}, & \text{if } 4 \leq z < 9 \\ 1, & \text{if } z \geq 9 \end{cases} \cdot \blacksquare$$

Example 2.2

Let X be a random variable with p.m.f.

$$f_X(x) = \begin{cases} \frac{|x|}{2550}, & \text{if } x \in \{\pm 1, \pm 2, \dots, \pm 50\} \\ 0, & \text{otherwise} \end{cases}.$$

Show that $Z = |X|$ is a random variable. Find its p.m.f., and distribution function.

Solution. As $h(x) = |x|, x \in \mathbb{R}$, is a continuous function and X is a random variable, using Remark 1.1 (i), $Z = |X|$ is a random variable. We have $S_X = \{\pm 1, \pm 2, \dots, \pm 50\}$ and $S_Z = \{1, 2, \dots, 50\}$. Moreover, for $z \in S_Z$,

$$P(\{Z = z\}) = P(\{|X| = z\}) = P(\{X \in \{-z, z\}\}) = \frac{|-z|}{2550} + \frac{|z|}{2550} = \frac{z}{1275}.$$

Therefore the p.m.f. of Z is

$$f_Z(z) = \begin{cases} \frac{z}{1275}, & \text{if } z \in \{1, 2, \dots, 50\} \\ 0, & \text{otherwise} \end{cases},$$

and the distribution function of Z is

$$F_Z(z) = \begin{cases} 0, & \text{if } z < 1 \\ \frac{1}{1275}, & \text{if } 1 \leq z < 2 \\ \frac{i(i+1)}{2550}, & \text{if } i \leq z < i+1, i = 2, 3, \dots, 49 \\ 1, & \text{if } z \geq 50 \end{cases} \quad \blacksquare$$

Example 2.3

Let X be a random variable with p.m.f.

$$f_X(x) = \begin{cases} \binom{n}{x} p^x (1-p)^{n-x}, & \text{if } x \in \{0, 1, \dots, n\} \\ 0, & \text{otherwise} \end{cases}$$

where n is a positive integer and $p \in (0,1)$. Show that $Y = n - X$ is a random variable. Find its p.m.f. and distribution function.

Solution. Note that $S_X = S_Y = \{0, 1, \dots, n\}$ and $h(x) = n - x, x \in \mathbb{R}$, is a continuous function. Therefore $Y = n - X$ is a random variable. For $y \in S_Y$

$$P(\{Y = y\}) = P(\{X = n - y\}) = \binom{n}{n-y} p^{n-y} (1-p)^y = \binom{n}{y} (1-p)^y p^{n-y}.$$

Thus the p.m.f. of Y is

$$f_Y(y) = \begin{cases} \binom{n}{y} (1-p)^y p^{n-y}, & \text{if } y \in \{0, 1, \dots, n\} \\ 0, & \text{otherwise} \end{cases}$$

and the distribution function of Y is

$$F_Y(y) = \begin{cases} 0, & \text{if } y < 0 \\ p^n, & \text{if } 0 \leq y < 1 \\ \sum_{j=0}^i \binom{n}{j} (1-p)^j p^{n-j}, & \text{if } i \leq y < i+1, i = 1, 2, \dots, n-1 \\ 1, & \text{if } y \geq n \end{cases} \quad \blacksquare$$

The following theorem deals with probability distribution of absolutely continuous type random variables.

Theorem 2.2

Let X be a random variable of absolutely continuous type with p.d.f. $f_X(\cdot)$ and support S_X . Let S_1, S_2, \dots, S_k , be open intervals in \mathbb{R} such that $S_i \cap S_j = \emptyset$, if $i \neq j$ and $\bigcup_{i=1}^k S_i = S_X$. Let $h: \mathbb{R} \rightarrow \mathbb{R}$ be a Borel function such that, on each S_i ($i = 1, \dots, k$), $h: S_i \rightarrow \mathbb{R}$ is strictly monotone and continuously differentiable with inverse function $h_i^{-1}(\cdot)$. Let $h(S_j) = \{h(x): x \in S_j\}$ so that $h(S_j)$ ($j = 1, \dots, k$) is an open interval in \mathbb{R} . Then the random variable $T = h(X)$ is of absolutely continuous type with p.d.f.

$$f_T(t) = \sum_{j=1}^k f_X(h_j^{-1}(t)) \left| \frac{d}{dt} h_j^{-1}(t) \right| I_{h(S_j)}(t).$$

Proof. We will provide an outline of the proof which may not be rigorous. Let $F_T(\cdot)$ be the distribution function of T . For $t \in \mathbb{R}$ and $\Delta > 0$,

$$\begin{aligned} \frac{F_T(t + \Delta) - F_T(t)}{\Delta} &= \frac{P(\{t < h(X) \leq t + \Delta\})}{\Delta} \\ &= \sum_{j=1}^k \frac{P(\{t < h(X) \leq t + \Delta, X \in S_j\})}{\Delta}. \end{aligned}$$

Fix $j \in \{1, \dots, k\}$. First suppose that $h_j(\cdot)$ is strictly decreasing on S_j . Note that $\{X \in S_j\} = \{h(X) \in h(S_j)\}$ and $h(S_j)$ is an open interval. Thus, for t belonging to the exterior of $h(S_j)$ and sufficiently small $\Delta > 0$, we have $P(\{t < h(X) \leq t + \Delta, X \in S_j\}) = 0$. Also, for $t \in h(S_j)$ and sufficiently small $\Delta > 0$,

$$P(\{t < h(X) \leq t + \Delta, X \in S_j\}) = P(\{h_j^{-1}(t + \Delta) \leq X < h_j^{-1}(t)\}).$$

Thus, for all $t \in \mathbb{R}$, we have

$$\begin{aligned} \frac{P(\{t < h(X) \leq t + \Delta, X \in S_j\})}{\Delta} &= \frac{P(\{h_j^{-1}(t + \Delta) \leq X < h_j^{-1}(t)\}) I_{h(S_j)}(t)}{\Delta} \\ &= \frac{1}{\Delta} \left[\int_{h_j^{-1}(t+\Delta)}^{h_j^{-1}(t)} f_X(z) dz \right] I_{h(S_j)}(t) \\ &\xrightarrow{\Delta \downarrow 0} -f_X(h_j^{-1}(t)) \left(\frac{d}{dt} h_j^{-1}(t) \right) I_{h(S_j)}(t). \end{aligned} \quad (2.1)$$

Similarly if h_j is strictly increasing on S_j then, for all $t \in \mathbb{R}$, we have

$$\begin{aligned}
\frac{P(\{t < h(X) \leq t + \Delta, X \in S_j\})}{\Delta} &= \frac{P(\{h_j^{-1}(t) < X \leq h_j^{-1}(t + \Delta)\}) I_{h(S_j)}(t)}{\Delta} \\
&= \frac{1}{\Delta} \left[\int_{h_j^{-1}(t)}^{h_j^{-1}(t+\Delta)} f_X(z) dz \right] I_{h(S_j)}(t) \\
&\xrightarrow{\Delta \downarrow 0} f_X(h_j^{-1}(t)) \left(\frac{d}{dt} h_j^{-1}(t) \right) I_{h(S_j)}(t). \tag{2.2}
\end{aligned}$$

Note that if h is strictly decreasing (increasing) on S_j then $\frac{d}{dt} h_j^{-1}(t) < (>) 0$ on S_j . Now on combining (2.1) and (2.2) we get, for all $t \in \mathbb{R}$,

$$\begin{aligned}
\frac{P(\{t < h(X) \leq t + \Delta, X \in S_j\})}{\Delta} &\xrightarrow{\Delta \downarrow 0} f_X(h_j^{-1}(t)) \left| \frac{d}{dt} h_j^{-1}(t) \right| I_{h(S_j)}(t), \\
\Rightarrow \frac{F_T(t + \Delta) - F_T(t)}{\Delta} &\xrightarrow{\Delta \downarrow 0} \sum_{j=1}^k f_X(h_j^{-1}(t)) \left| \frac{d}{dt} h_j^{-1}(t) \right| I_{h(S_j)}(t).
\end{aligned}$$

Similarly one can show that, for all $t \in \mathbb{R}$,

$$\lim_{\Delta \downarrow 0} \frac{F_T(t + \Delta) - F_T(t)}{\Delta} = \sum_{j=1}^k f_X(h_j^{-1}(t)) \left| \frac{d}{dt} h_j^{-1}(t) \right| I_{h(S_j)}(t). \tag{2.3}$$

It follows that the distribution function of T is differentiable everywhere on \mathbb{R} except possibly at a finite number of points (on boundaries of intervals $h(S_1), \dots, h(S_k)$ of S_T). Now the result follows from Remark 4.2 (vii) of Module 2 and using (2.3). ■

The following corollary to the above theorem is immediate.

Corollary 2.2

Let X be a random variable of absolutely continuous type with p.d.f. $f_X(\cdot)$ and support S_X . Suppose that S_X is a finite union of disjoint open intervals in \mathbb{R} and let $h: \mathbb{R} \rightarrow \mathbb{R}$ be a Borel function such that h is differentiable and strictly monotone on S_X (i.e., either $h'(x) < 0, \forall x \in S_X$ or $h'(x) > 0, \forall x \in S_X$). Let $S_T = \{h(x): x \in S_X\}$. Then $T = h(X)$ is a random variable of absolutely continuous type with p.d.f.

$$f_T(t) = \begin{cases} f_X(h^{-1}(t)) \left| \frac{d}{dt} h^{-1}(t) \right|, & \text{if } t \in S_T \\ 0, & \text{otherwise} \end{cases} \quad \blacksquare$$

It may be worth mentioning here that, in view of Remark 4.2 (vii) of Module 2, Theorem 2.2 and Corollary 2.2 can be applied even in situations where the function h is differentiable everywhere on S_X except possibly at a finite number of points.

Example 2.4

Let X be random variable with p.d.f.

$$f_X(x) = \begin{cases} e^{-x}, & \text{if } x > 0 \\ 0, & \text{otherwise} \end{cases},$$

and let $T = X^2$

- (i) Show that T is a random variable of absolutely continuous type;
- (ii) Find the distribution function of T and hence find its p.d.f.;
- (iii) Find the p.d.f. of T directly (i.e., without finding the distribution function of T).

Solution. (i) and (iii). Clearly $T = X^2$ is a random variable (being a continuous function of random variable X). We have $S_X = S_T = (0, \infty)$. Also $h(x) = x^2, x \in S_X$, is strictly increasing on S_X with inverse function $h^{-1}(x) = \sqrt{x}, x \in S_T$. Using Corollary 2.1 it follows that $T = X^2$ is a random variable of absolutely continuous type with p.d.f.

$$\begin{aligned} f_T(t) &= \begin{cases} f_X(\sqrt{t}) \left| \frac{d}{dt}(\sqrt{t}) \right|, & \text{if } t > 0 \\ 0, & \text{otherwise} \end{cases} \\ &= \begin{cases} \frac{e^{-\sqrt{t}}}{2\sqrt{t}}, & \text{if } t > 0 \\ 0, & \text{otherwise} \end{cases}. \end{aligned}$$

(ii) We have $F_T(t) = P(\{X^2 \leq t\}), t \in \mathbb{R}$. Clearly, for $t < 0, F_T(t) = 0$. For $t \geq 0$,

$$\begin{aligned} F_T(t) &= P(\{-\sqrt{t} \leq X \leq \sqrt{t}\}) \\ &= \int_{-\sqrt{t}}^{\sqrt{t}} f_X(x) dx \\ &= \int_0^{\sqrt{t}} e^{-x} dx \\ &= 1 - e^{-\sqrt{t}}. \end{aligned}$$

Therefore the distribution function of T is

$$F_T(t) = \begin{cases} 0, & \text{if } t < 0 \\ 1 - e^{-\sqrt{t}}, & \text{if } t \geq 0 \end{cases}$$

Clearly F_T is differentiable everywhere except at $t = 0$. Therefore, using Remark 4.2 (vii) of Module 2, we conclude that the random variable T is of absolutely continuous type with p.d.f. $f_T(t) = F_T'(t)$, if $t \neq 0$. At $t = 0$ we may assign any arbitrary non-negative value to $f_T(0)$. Thus a p.d.f. of T is

$$f_T(t) = \begin{cases} \frac{e^{-\sqrt{t}}}{2\sqrt{t}}, & \text{if } t > 0 \\ 0, & \text{otherwise} \end{cases} \quad \blacksquare$$

Example 2.5

Let X be a random variable with p.d.f.

$$f_X(x) = \begin{cases} \frac{|x|}{2}, & \text{if } -1 < x < 1 \\ \frac{x}{3}, & \text{if } 1 \leq x < 2 \\ 0, & \text{otherwise} \end{cases} \quad ,$$

and let $T = X^2$

- (i) Show that T is a random variable of absolutely continuous type;
- (ii) Find the distribution function of T and hence find its p.d.f;
- (iii) Find the p.d.f. of T directly (i.e., without finding the distribution function of T).

Solution. (i) and (iii). Clearly $T = X^2$ is a random variable (being a continuous function of random variable X). We have $S_X = (-1, 0) \cup (0, 2) = S_1 \cup S_2$, say. Also $h(x) = x^2, x \in S_X$, is strictly decreasing in $S_1 = (-1, 0)$ with inverse function $h_1^{-1}(t) = -\sqrt{t}$; $h(x) = x^2, x \in S_X$, is strictly increasing in $S_2 = (0, 2)$, with inverse function $h_2^{-1}(t) = \sqrt{t}$; $h(S_1) = (0, 1)$ and $h(S_2) = (0, 4)$. Using Theorem 2.2 it follows that $T = X^2$ is a random variable of absolutely continuous type with p.d.f.

$$\begin{aligned} f_T(t) &= f_X(-\sqrt{t}) \left| \frac{d}{dt}(-\sqrt{t}) \right| I_{(0,1)}^{(t)} + f_X(\sqrt{t}) \left| \frac{d}{dt}(\sqrt{t}) \right| I_{(0,4)}^{(t)} \\ &= \begin{cases} \frac{1}{2}, & \text{if } 0 < t < 1 \\ \frac{1}{6}, & \text{if } 1 < t < 4 \\ 0, & \text{otherwise} \end{cases} \end{aligned}$$

(ii) We have $F_T(t) = P(\{X^2 \leq t\}), t \in \mathbb{R}$. Since $P(\{X \in (-1, 2)\}) = 1$, we have $P(\{T \in (0, 4)\}) = 1$.

Therefore, for $t < 0$, $F_T(t) = P(\{T \leq t\}) = 0$ and, for $t \geq 4$, $F_T(t) = P(\{T \leq t\}) = 1$. For $t \in [0, 4)$, we have

$$\begin{aligned} F_T(t) &= P(\{-\sqrt{t} \leq X \leq \sqrt{t}\}) \\ &= \int_{-\sqrt{t}}^{\sqrt{t}} f_X(x) dx \\ &= \begin{cases} \int_{-\sqrt{t}}^{\sqrt{t}} \frac{|x|}{2} dx, & \text{if } 0 \leq t < 1 \\ \int_{-1}^1 \frac{|x|}{2} dx + \int_1^{\sqrt{t}} \frac{x}{3} dx, & \text{if } 1 \leq t < 4 \end{cases}. \end{aligned}$$

Therefore, the distribution function of T is

$$F_T(t) = \begin{cases} 0, & \text{if } t < 0 \\ \frac{t}{2}, & \text{if } 0 \leq t < 1 \\ \frac{t+2}{6}, & \text{if } 1 \leq t < 4 \\ 1, & \text{if } t \geq 4 \end{cases}.$$

Clearly F_T is differentiable everywhere except at points 0, 1 and 4. Using Remark 4.2 (vii) of Module 2 it follows that the random variable T is of absolutely continuous type with a p.d.f.

$$f_T(t) = \begin{cases} \frac{1}{2}, & \text{if } 0 < t < 1 \\ \frac{1}{6}, & \text{if } 1 < t < 4 \\ 0, & \text{otherwise} \end{cases}. \blacksquare$$

Note that a Borel function of a discrete type random variable is a random variable of discrete type (see Theorem 1.1). Theorem 2.2 provides sufficient conditions under which a Borel function of an absolutely continuous type random variable is of absolutely continuous type. The following example illustrates that, in general, a Borel function of an absolutely continuous type random variable may not be of absolutely continuous type.

Example 2.6

Let X be a random variable of absolutely continuous type with p.d.f.

$$f_X(x) = \begin{cases} e^{-x}, & \text{if } x > 0 \\ 0, & \text{otherwise} \end{cases}$$

and let $T = [X]$, where, for $x \in \mathbb{R}$, $[x]$ denotes the largest integer not exceeding x . Show that T is a random variable of discrete type and find its p.m.f.

Solution. For $a \in \mathbb{R}$, we have

$$T^{-1}((-\infty, a]) = (-\infty, [a] + 1) \in \mathcal{B}_1.$$

It follows that T is a random variable. Also $S_X = (0, \infty)$. Since $P(\{X \in S_X\}) = 1$, we have $P(T \in \{0, 1, 2, \dots\}) = 1$. Also, for $i \in \{0, 1, 2, \dots\}$.

$$\begin{aligned} P(\{T = i\}) &= P(\{i \leq X < i + 1\}) \\ &= \int_i^{i+1} f_X(x) dx \\ &= \int_i^{i+1} e^{-x} dx \\ &= (1 - e^{-1})e^{-i} \\ &> 0. \end{aligned}$$

Consequently the random variable T is of discrete type with support $S_T = \{0, 1, 2, \dots\}$ and p.m.f.

$$f_T(t) = P(\{T = t\}) = \begin{cases} (1 - e^{-1})e^{-t}, & \text{if } t \in \{0, 1, 2, \dots\} \\ 0, & \text{otherwise} \end{cases}. \blacksquare$$