

1 Functions of two or more than one variable

In a 2-dimensional space a point is represented by an ordered pair (x, y) and the set of all such points constitutes a 2-dimensional Euclidean space $\mathbb{R}^2$. For $\mathbb{R}^n$, a point will be of form $(x_1, x_2, \dots, x_n)$. If in 2D space the variables x and y are independent such that a variable $z = f(x, y)$ is a function of x and y . Here z is dependent variable. The subset D of $\mathbb{R}^2$ is the set of points for which the function $f(x, y)$ is defined, then D is called domain of f .

Example 1.1. $z = \frac{xy}{x^2 - y^2}$. Here z is not defined for $x = y$.

$$D = \{(x, y) : x \neq y\} = \{(x, y) \in \mathbb{R}^2 : x \neq y\} = \{x \in \mathbb{R}, y \in \mathbb{R} : x \neq y\}$$

Example 1.2. $z = \sqrt{1 - x - y}$. Here $D = \{(x, y) \in \mathbb{R}^2 : x + y \leq 1\}$

Limit of $f(x, y)$. Let D be a subset of $\mathbb{R}^2$ which is domain of definition of a function f . Let (a, b) be a point in D . Let $(x, y) \in D$ and $(x, y) \rightarrow (a, b)$ in any manner. f is said to tend to a limit A if $\forall \varepsilon > 0 \exists \delta > 0$ such that $|f(x, y) - A| < \varepsilon$, whenever $(x, y) \in$ a nbd of the point (a, b) . We may choose the nbd as

$$N_{(a,b)} = \{(x, y) \in \mathbb{R}^2 : 0 < |x - a| < \delta, 0 < |y - b| < \delta\} \quad \text{or,}$$

$$N_{(a,b)} = \{(x, y) \in \mathbb{R}^2 : (x - a)^2 + (y - b)^2 < \delta^2\}$$

Then the limit, $\lim_{(x,y) \rightarrow (a,b)} f(x, y) = A$ is called double limit or simultaneous limit.

Iterated or Repeated Limit. Let N be an nbd of (a, b) and $f(x, y)$ is defined on N . For a fixed value of y , $\lim_{x \rightarrow a} f(x, y)$, if exists, will involve y . Therefore, $\lim_{x \rightarrow a} f(x, y)$, will be different for different values of y . That is $\lim_{x \rightarrow a} f(x, y) = \phi(y)$. Again if $\lim_{y \rightarrow b} \phi(y)$, exists and equal to A , then we write $\lim_{y \rightarrow b} \lim_{x \rightarrow a} f(x, y) = A$.

Example 1.3. Find the limit $\lim_{x \rightarrow a} f(x, y)$, where $f(x, y) = xy \frac{x^2 - y^2}{x^2 + y^2}$.

Sol. Let us take $x = r \cos \theta, y = r \sin \theta$. Then $x^2 + y^2 = r^2$ and as $(x, y) \rightarrow (0, 0), r \rightarrow 0$
 $f(x, y) = r^2 \sin \theta \cos \theta \cos 2\theta$,

$$\left| xy \frac{x^2 - y^2}{x^2 + y^2} \right| = \left| \frac{r^2}{4} \sin 4\theta \right| \leq \frac{r^2}{4} < \varepsilon, \quad \text{if } r^2 < 4\varepsilon \quad \text{or} \quad 0 < x^2 + y^2 < \delta^2.$$

So, $\lim_{(x,y) \rightarrow (0,0)} f(x, y) = 0$ (using definition).

Example 1.4. Let, $f(x, y) = \begin{cases} x \sin \frac{1}{y} + \frac{x^2 - y^2}{x^2 + y^2} & y \neq 0 \\ 0 & y = 0 \end{cases}$

Show that $\lim_{y \rightarrow 0} \lim_{x \rightarrow 0} f(x, y)$ exists but neither of $\lim_{(x,y) \rightarrow (0,0)} f(x, y)$ nor $\lim_{x \rightarrow 0} \lim_{y \rightarrow 0} f(x, y)$ exists.

Sol. We shall show that $\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 - y^2}{x^2 + y^2}$ does not exist. Since $(x, y) \rightarrow (0, 0)$, let, $y = mx$ be the path.

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 - y^2}{x^2 + y^2} = \lim_{(x,y) \rightarrow (0,0)} \frac{x^2 - m^2 x^2}{x^2 + m^2 x^2} = \lim_{x \rightarrow 0} \frac{1 - m^2}{1 + m^2} = \frac{1 - m^2}{1 + m^2} \quad (x \neq 0)$$

So, the limit is path dependant, i.e., the limit is different for different values of m .

Again, $\lim_{y \rightarrow 0} \sin \frac{1}{y}$ does not exist, then $\lim_{x \rightarrow 0} \lim_{y \rightarrow 0} x \sin \frac{1}{y}$ does not exist therefore, $\lim_{x \rightarrow 0} \lim_{y \rightarrow 0} f(x, y)$ does not exist.

Now, $\lim_{x \rightarrow 0} x \sin \frac{1}{y} = 0$ and $\lim_{x \rightarrow a} f(x, y) = -1$, then $\lim_{y \rightarrow 0} (0 - 1) = -1$
 ,i.e., $\lim_{y \rightarrow 0} \lim_{x \rightarrow 0} (x \sin \frac{1}{y} + \frac{x^2 - y^2}{x^2 + y^2}) = -1$

Partial Derivatives for functions of two or more variables. $D \subset \mathbb{R}$ is usual domain of f and let $P(a, b) \in D$ be arbitrary. Let N be an nbd and $Q(a + h, b + k) \in N$. Then if α be an angle that the line joining $\vec{PQ}$ makes with positive x -axis, then the direction cosines of $\vec{PQ}$ are $l = \cos \alpha, m = \sin \alpha$. If $\rho = \sqrt{h^2 + k^2}$. Then $h = \rho \cos \alpha, k = \rho \sin \alpha$ and $\rho \rightarrow 0$ as $Q \rightarrow P$. Now if,

$$\lim_{\rho \rightarrow 0} \frac{f(a + \rho \cos \alpha, b + \rho \sin \alpha) - f(a, b)}{\rho} \quad \text{exists.}$$

Then, it is called derivative of $f(x, y)$ at (a, b) in direction of α and it is denoted by $D_\alpha f(a, b)$. If $\alpha = 0$, then derivative is denoted by $\frac{\partial f}{\partial x}$ and if $\alpha = \frac{\pi}{2}$, then the derivative is denoted by $\frac{\partial f}{\partial y}$. These derivatives are called partial derivatives.

Continuity of $f(x, y)$. Let $N \subset D \subset \mathbb{R}^2$, where D is the domain of and N is the nbd of a point $(a, b) \in D$. Then f is said to be continuous at (a, b) if $\lim_{(x,y) \rightarrow (a,b)} f(x, y) = f(a, b)$.

Equivalently, $\forall \varepsilon > 0 \exists \delta > 0$ such that $|f(x, y) - f(a, b)| < \varepsilon$ whenever $|x - a| < \delta$ and $|y - b| < \delta$ or $(x - a)^2 + (y - b)^2 < \delta^2$
 or,

$$\lim_{(h,k) \rightarrow (0,0)} f(a + h, b + k) = f(a, b) \quad \text{where } (a + h, b + k) \in D.$$

Example 1.5. Verify the continuity of $f(x, y) = \begin{cases} xy \frac{x^2 - y^2}{x^2 + y^2} & (x, y) \neq (0, 0) \\ 0 & (x, y) = (0, 0) \end{cases}$ at $(0, 0)$.

Sol. Here,

$$|f(x, y) - f(0, 0)| = |xy \frac{x^2 - y^2}{x^2 + y^2} - 0| = |xy \frac{x^2 - y^2}{x^2 + y^2}| = |x||y| \frac{|x^2 - y^2|}{x^2 + y^2} \leq |x||y| \leq \frac{x^2 + y^2}{2} < \varepsilon,$$

if $x^2 + y^2 < \delta^2 (= 2\varepsilon) \Rightarrow f$ is continuous at $(0, 0)$.

Example 1.6. Verify the continuity of $f(x, y) = \begin{cases} x \sin \frac{1}{y} + y \sin \frac{1}{x} & xy \neq 0 \\ 0 & xy = 0 \end{cases}$ at $(0, 0)$.

Sol. Here,

$$|f(x, y) - f(0, 0)| = |x \sin \frac{1}{y} + y \sin \frac{1}{x} - 0| \leq |x| \sin \frac{1}{y} + |y| \sin \frac{1}{x} \leq |x| + |y| < \varepsilon$$

if $|x - 0| < \delta$, and $|y - 0| < \delta (= \frac{\varepsilon}{2}) \Rightarrow f$ is continuous at $(0, 0)$.

Example 1.7. Verify the continuity of $f(x, y) = \begin{cases} \frac{xy}{x^2 + y^2} & x^2 + y^2 \neq 0 \\ 0 & x^2 + y^2 = 0 \end{cases}$ at $(0, 0)$.

Partial Derivatives (Computations). If $\lim_{h \rightarrow 0} \frac{f(x+h, y) - f(x, y)}{h}$ exists, then it is denoted by $\frac{\partial f}{\partial x}$ or f_x similarly, if $\lim_{k \rightarrow 0} \frac{f(x, y+k) - f(x, y)}{k}$ exists, then it is denoted by $\frac{\partial f}{\partial y}$ or f_y .

Example 1.8. Let, $f(x, y) = x^2y + 3xy$. Find f_x, f_y .

Sol. $f_x = 2xy + 3y$ and $f_y = x^2 + 3x$.

Let $z = f(x, y)$ then,

$$\begin{aligned} z_x &= \frac{\partial z}{\partial x} = \frac{\partial f}{\partial x}, z_y = \frac{\partial z}{\partial y} = \frac{\partial f}{\partial y}, \\ z_{xx} &= \frac{\partial}{\partial x} \left(\frac{\partial z}{\partial x} \right) = \frac{\partial^2 z}{\partial x^2}, z_{yy} = \frac{\partial}{\partial y} \left(\frac{\partial z}{\partial y} \right) = \frac{\partial^2 z}{\partial y^2}, z_{yx} = \frac{\partial}{\partial y} \left(\frac{\partial z}{\partial x} \right) = \frac{\partial^2 z}{\partial y \partial x} \dots, \\ \frac{\partial}{\partial y} \left(\frac{\partial^{n-1}}{\partial y^{n-1}} z \right) &= \frac{\partial^n z}{\partial y^n}, \frac{\partial}{\partial x} \left(\frac{\partial^{n-1}}{\partial x^{n-1}} z \right) = \frac{\partial^n z}{\partial x^n}. \end{aligned}$$

Example 1.9. $z = f(x, y) = x^3 + 3xy^4 - 3x^4y^3$. Find $z_x, z_y, z_{xx}, z_{yy}, z_{yx}, z_{xy}$.

Sol.

$$\begin{aligned} z_x &= 3x^2 + 3y^4 - 12x^3y^3, \\ z_y &= 12xy^3 - 9x^4y^2, \\ z_{xx} &= 6x - 36x^2y^3, \\ z_{yy} &= 36xy^2 - 24x^4y, \\ z_{yx} &= 12y^3 - 36x^3y^2, \\ z_{xy} &= 12y^3 - 36x^3y^2 \end{aligned}$$

Taylor's Series: The Taylor's series of a function $f(x, y)$ about a point (x_0, y_0) provide an approximation of a function in the nbd of (x_0, y_0) and

$$\begin{aligned} z(x, y) &= z(x_0, y_0) + \frac{\partial z}{\partial x} \Big|_{(x_0, y_0)} (x - x_0) + \frac{\partial z}{\partial y} \Big|_{(x_0, y_0)} (y - y_0) + \frac{1}{2} \left(\frac{\partial^2 z}{\partial x^2} \Big|_{(x_0, y_0)} (x - x_0)^2 \right. \\ &\quad \left. + \frac{\partial^2 z}{\partial y^2} \Big|_{(x_0, y_0)} (y - y_0)^2 + 2 \frac{\partial^2 z}{\partial x \partial y} \Big|_{(x_0, y_0)} (x - x_0)(y - y_0) \right) + \dots \end{aligned}$$

Example 1.10. Let us take the function, $z(x, y) = x^3 + 3y - y^3 - 3x$. Find $z_x, z_y, z_{xx}, z_{yy}, z_{yx}, z_{xy}$.

Sol. $z_x = 3x^2 - 3, z_y = 3 - 3y^2, z_{xx} = 6x, z_{yy} = -6y, z_{yx} = 0, z_{xy} = 0$.

2 Maxima and Minima

Stationary Points: Maxima and Minima:

If $f'(x) = 0$ at $x = x_0$ and $f''(x) \Big|_{x=x_0} > 0$ then f attains minimum value at $x = x_0$. Similarly, if $f'(x) = 0$ at $x = x_0$ and $f''(x) \Big|_{x=x_0} < 0$ then f attains maximum value at $x = x_0$.

If f is a function of two variables, $\frac{\partial f}{\partial x} = 0, \frac{\partial f}{\partial y} = 0 \Rightarrow x = x_0, y = y_0$. Then we will calculate second order derivatives of it at point (x_0, y_0) .

Let, $z_{xx}|_{(x_0, y_0)} = A$, $z_{yy}|_{(x_0, y_0)} = B$, $z_{xy}|_{(x_0, y_0)} = C$. Compute the discriminant: $D = BC - A^2$

Case I. If $D > 0$ and $\frac{\partial^2 z}{\partial x^2}|_{(x_0, y_0)} > 0$ or $\frac{\partial^2 z}{\partial y^2}|_{(x_0, y_0)} > 0$, then z attains local minima at point (x_0, y_0) .

Case II. If $D > 0$ and $\frac{\partial^2 z}{\partial x^2}|_{(x_0, y_0)} < 0$ or $\frac{\partial^2 z}{\partial y^2}|_{(x_0, y_0)} < 0$, then z attains local maxima at point (x_0, y_0) .

Case III. If $D < 0$, then (x_0, y_0) is a saddle point.

Example 2.1. Find the maxima and minima of $f(x) = x^3 - 4x^2$.

Sol. Given

$$\begin{aligned} f(x) &= x^3 - 4x^2 \\ \Rightarrow f'(x) &= 3x^2 - 8x \end{aligned}$$

For extrema/ extremum,

$$f'(x) = 0 \Rightarrow 3x^2 - 8x = 0 \Rightarrow x = 0, x = \frac{8}{3}.$$

Again,

$$\begin{aligned} f''(x) &= 6x - 8 \\ f''(0) &= -8 < 0 \\ f''\left(\frac{8}{3}\right) &= 6 \times \frac{8}{3} - 8 = 16 - 8 = 8 > 0. \end{aligned}$$

At, $x = 0$, $f''(x) < 0$ which implies that the function attains its maximum value at $x = 0$.
At, $x = \frac{8}{3}$, $f''(x) > 0$, which implies that the function attains its minimum value at $x = \frac{8}{3}$.
So, the maximum value of

$$\begin{aligned} f &= f(x)|_{x=0} \\ &= 0 \end{aligned}$$

And the minimum value of

$$\begin{aligned} f &= f(x)|_{x=\frac{8}{3}} \\ &= x^3 - 4x^2|_{x=\frac{8}{3}} \\ &= \frac{8^3}{27} - \frac{4 \cdot 8^2}{9} \\ &= \frac{8^2}{9} \times \left[\frac{8}{3} - 4\right] \\ &= \frac{8^2}{9} \times \left(-\frac{4}{3}\right) \\ &= -\frac{256}{27} \end{aligned}$$

Example 2.2. Find the maximum and minimum values of the function,

$$f(x, y) = x^3 - y^3 + 3y - 3x$$

Sol. Here,

$$\begin{aligned}f_x(x, y) &= 3x^2 - 3 \\f_y(x, y) &= -3y^2 + 3\end{aligned}$$

Suppose the extremum values are attained at the point (x_0, y_0) . Then for the extremum,

$$\begin{aligned}f_x(x_0, y_0) = 0 &\Rightarrow 3x_0^2 - 3 = 0 \Rightarrow x_0^2 = 1 \Rightarrow x_0 = \pm 1 \\f_y(x_0, y_0) = 0 &\Rightarrow -3y_0^2 + 3 = 0 \Rightarrow y_0^2 = 1 \Rightarrow y_0 = \pm 1\end{aligned}$$

The extremum points are $P_1(1, 1)$, $P_2(1, -1)$, $P_3(-1, 1)$, $P_4(-1, -1)$. For maximum and minimum values, we need the 2nd derivatives and the discriminant D,

$$\begin{aligned}f_{xx} &= 6x, f_{yy} = -6y, f_{xy} = 0, f_{yx} = 0 \\D &= AB - C^2 = 6x \times (-6y) - 0 = -36xy\end{aligned}$$

Now,

$$\begin{aligned}D\Big|_{P_1(1,1)} &= -36 \times 1 \times 1 = -36 < 0, A = 6, B = -6, C = 0 \rightarrow \text{Saddle Point} \\D\Big|_{P_2(1,-1)} &= 36 > 0, A = 6, B = 6, C = 0 \rightarrow \text{Minimum Value} \\D\Big|_{P_3(-1,1)} &= 36 > 0, A = -6, B = -6, C = 0 \rightarrow \text{Maximum Value} \\D\Big|_{P_4(-1,-1)} &= -36 < 0, A = -6, B = 6, C = 0 \rightarrow \text{Saddle Point} \\f(x, y)\Big|_{(1,-1)} &= 8, f(x, y)\Big|_{(-1,1)} = 4\end{aligned}$$

Example 2.3. For the function

$$f(x, y) = x^3 - y^3 + 3y - 3x$$

Find out the Taylor's series expansion about the point $P(2, 1)$.

Sol.

$$\begin{aligned}f(x, y) &= x^3 - y^3 + 3y - 3x \Rightarrow f(x, y)\Big|_{P(2,1)} = 4 \\f_x(x, y) &= 3x^2 - 3 \Rightarrow f_x(x, y)\Big|_{P(2,1)} = 9 \\f_y(x, y) &= -3y^2 + 3 \Rightarrow f_y(x, y)\Big|_{P(2,1)} = 0 \\f_{xx}(x, y) &= 6x \Rightarrow f_{xx}(x, y)\Big|_{P(2,1)} = 12 \\f_{yy}(x, y) &= -6y \Rightarrow f_{yy}(x, y)\Big|_{P(2,1)} = -6 \\f_{xy}(x, y) &= 0 \Rightarrow f_{xy}(x, y)\Big|_{P(2,1)} = 0\end{aligned}$$

By Taylor's series,

$$\begin{aligned}f(x, y) &= f((2, 1)) + \frac{\partial f}{\partial x}\Big|_{(2,1)}(x - 2) + \frac{\partial f}{\partial y}\Big|_{(2,1)}(y - 1) + \frac{1}{2}\left(\frac{\partial^2 z}{\partial x^2}\Big|_{(2,1)}(x - 2)^2 + \frac{\partial^2 z}{\partial y^2}\Big|_{(2,1)}(y - 1)^2\right. \\&\quad \left.+ 2\frac{\partial^2 z}{\partial x \partial y}\Big|_{(2,1)}(x - 2)(y - 1)\right) + \dots \\&= 4 + 9 \times (x - 2) + 0 \times (y - 1) + 12 \times \frac{(x - 2)^2}{2} + (-6) \times \frac{(y - 1)^2}{2} + 0 \times (x - 2)(y - 1) + \dots \\&= 4 + 9(x - 2) + 6(x - 2)^2 - 3(y - 1)^2 + \dots\end{aligned}$$

Example 2.4. If $u = \tan^{-1} \frac{x^3+y^3}{x-y}$, prove that $x^2 u_{xx} + 2xy u_{xy} + y^2 u_{yy} = (1 - 4 \sin^2 u) \sin 2u$

Sol. Let, $v = \frac{x^3+y^3}{x-y} = \tan u$. Here, v is a homogeneous function of x and y of degree 2. Then by Euler's theorem,

$$\begin{aligned} x \frac{\partial v}{\partial x} + y \frac{\partial v}{\partial y} &= 2v = 2 \tan u \\ \Rightarrow x \frac{\partial v}{\partial u} \frac{\partial u}{\partial x} + y \frac{\partial v}{\partial u} \frac{\partial u}{\partial y} &= 2v = 2 \tan u \\ \Rightarrow \left(x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} \right) \sec^2 u &= 2 \tan u \\ \Rightarrow x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} &= \sin 2u \end{aligned}$$

Differentiating with respect to x and y respectively,

$$\frac{\partial u}{\partial x} + x \frac{\partial^2 u}{\partial x^2} + y \frac{\partial^2 u}{\partial x \partial y} = 2 \cos 2u \frac{\partial u}{\partial x} \quad (2.1)$$

$$x \frac{\partial^2 u}{\partial y \partial x} + \frac{\partial u}{\partial y} + y \frac{\partial^2 u}{\partial y^2} = 2 \cos 2u \frac{\partial u}{\partial y} \quad (2.2)$$

Now $2.1 \times x + (2.2) \times y$ we get,

$$\begin{aligned} x^2 u_{xx} + 2xy u_{xy} + y^2 u_{yy} + \sin 2u &= 2 \cos 2u \sin 2u \\ &= \sin 2u (2 \cos 2u - 1) \\ &= (1 - 4 \sin^2 u) \sin 2u \end{aligned}$$

3 Chapter 2: Integral Calculus

$$I = \int_a^b f(x) dx$$

If the integrand of the integral becomes infinite in the interval $[a, b]$ or f has points of infinite discontinuity or when both or either one of the limit points/ end points $[a, b]$, is infinite then the integral I is called improper integral.

Example 3.1. $\int_0^1 \frac{1}{x^2} dx, \int_1^2 \frac{1}{(2-x)(1-x)} dx, \int_0^1 \frac{1}{(x-\frac{1}{2})^2} dx, \int_0^\infty e^x dx, \int_{-\infty}^0 \frac{1}{e^x} dx$

Convergence at the left end point.

$$I = \int_a^b f(x) dx$$

Let, at $x = a$, $f(x)$ has infinite discontinuity. In this case, we first find the value of $\int_{a+\varepsilon}^b f(x) dx$ then pass the limit $\varepsilon \rightarrow 0^+$ i.e,

$$I = \lim_{\varepsilon \rightarrow 0^+} \int_{a+\varepsilon}^b f(x) dx$$

Convergence at the right end point.

$$I = \int_a^b f(x) dx$$

Let, at $x = b$, $f(x)$ has infinite discontinuity. In this case, we first find the value of $\int_a^{b+\varepsilon} f(x) dx$ then pass the limit $\varepsilon \rightarrow 0^-$ i.e,

$$\begin{aligned} I &= \lim_{\varepsilon \rightarrow 0^-} \int_a^{b+\varepsilon} f(x) dx \\ &= \lim_{\varepsilon \rightarrow 0^+} \int_a^{b-\varepsilon} f(x) dx \end{aligned}$$

Convergence inside the interval. Let at $x = c \in (a, b)$, $f(x)$ has an infinite discontinuity. In this case, we first separate $I = \int_a^c f(x) dx + \int_c^b f(x) dx$, then we do the same as before

$$\begin{aligned} I &= \int_a^b f(x) dx \\ &= \int_a^c f(x) dx + \int_c^b f(x) dx \\ &= \lim_{\varepsilon \rightarrow 0^-} \int_a^{c+\varepsilon} f(x) dx + \lim_{\varepsilon \rightarrow 0^+} \int_{c-\varepsilon}^b f(x) dx \end{aligned}$$

Convergence at $+\infty, -\infty$.

$$\begin{aligned} I &= \int_a^\infty f(x) dx \\ &= \lim_{t \rightarrow \infty} \int_a^t f(x) dx \\ I_1 &= \int_{-\infty}^b f(x) dx \\ &= \lim_{t \rightarrow -\infty} \int_t^b f(x) dx \\ I_2 &= \int_{-\infty}^\infty f(x) dx \\ &= \int_{-\infty}^c f(x) dx + \int_c^\infty f(x) dx \end{aligned}$$

Example 3.2. $I = \int_0^1 \frac{1}{\sqrt{1-x^2}} dx$

Sol. At $x = 1$, there is an infinite discontinuity.

$$\begin{aligned} I &= \lim_{\varepsilon \rightarrow 0^+} \int_0^{1-\varepsilon} \frac{1}{\sqrt{1-x^2}} dx \\ &= \lim_{\varepsilon \rightarrow 0^+} \left[\sin^{-1} x \right]_0^{1-\varepsilon} \\ &= \lim_{\varepsilon \rightarrow 0^+} \sin^{-1}(1-\varepsilon) \\ &= \sin^{-1} 1 \\ &= \frac{\pi}{2} \end{aligned}$$

Example 3.3. $I = \int_0^{\pi} \frac{1}{\sin x} dx$

Hint.

$$\begin{aligned} I &= \int_0^{\pi} \frac{1}{\sin x} dx \\ &= \int_0^{\frac{\pi}{2}} \frac{1}{\sin x} dx + \int_{\frac{\pi}{2}}^{\pi} \frac{1}{\sin x} dx \end{aligned}$$

Example 3.4. $I = \int_0^{\infty} \frac{1}{1+x^2} dx$

Sol.

$$\begin{aligned} I &= \int_0^{\infty} \frac{1}{1+x^2} dx = \lim_{t \rightarrow \infty} \int_0^t \frac{1}{1+x^2} dx = \lim_{t \rightarrow \infty} \left[\tan^{-1} x \right]_0^t \\ &= \lim_{t \rightarrow \infty} \tan^{-1} t \\ &= \frac{\pi}{2} \end{aligned}$$

Example 3.5. $I = \int_a^b \frac{1}{(x-a)^n} dx$

Sol.

$$\begin{aligned} I &= \int_a^b \frac{1}{(x-a)^n} dx \\ &= \lim_{\varepsilon \rightarrow 0^+} \int_{a+\varepsilon}^b \frac{1}{(x-a)^n} dx \\ &= \lim_{\varepsilon \rightarrow 0^+} \left[\frac{1}{1-n} (x-a)^{(1-n)} \right]_{a+\varepsilon}^b \\ &= \lim_{\varepsilon \rightarrow 0^+} \frac{1}{1-n} \left(\frac{1}{(b-a)^{(n-1)}} - \frac{1}{\varepsilon^{n-1}} \right) \end{aligned}$$

It is convergent if $n < 1$ and divergent if $n \geq 1$.